

We've got to stop  
meeting like this.



Worried about how many times you excuse yourself?

You can do something about your bladder control problem.  
Ask your doctor about DETROL LA.

Have strong, sudden urges to go to the bathroom, and maybe even accidents? Go more than 8 times a day? If so, you may have a medical condition called overactive bladder. But with DETROL LA, you can help reduce the symptoms so you can worry less about your bladder.

One DETROL LA works 24 hours, all day and all night.\* It helps calm the bladder muscle that causes those strong, sudden urges. Just imagine what a calmer bladder can do for you.

1 DETROL LA 24 HOURS

**Detrol<sup>®</sup> LA**

tolterodine tartrate  
extended release capsules

*Helps keep an overactive  
bladder quieter.<sup>™</sup>*

If you have certain stomach problems, glaucoma, or trouble getting urine to pass, you shouldn't take DETROL LA. The most common side effects are dry mouth, headache, constipation, and abdominal pain.

DETROL LA, like all medicines, has benefits and risks. There may be other options. Ask your doctor if DETROL LA is right for you. For more information, visit [DETROLLA.com](http://DETROLLA.com).

\*Results may vary. Please see important product information on back. © 2007 Pfizer Inc. All rights reserved. April 2007 DD275001E

Uninsured? Need help paying for medicine? Pfizer has programs that can help, no matter your age or income. You may even qualify for free Pfizer medicines. Call 1-866-706-2400. Or visit [www.pfizerhelpfulanswers.com](http://www.pfizerhelpfulanswers.com).

Pfizer  
**helpful  
answers<sup>™</sup>**

# IMPORTANT FACTS

**Detrol<sup>®</sup> LA**  
tolterodine tartrate  
extended release capsules

(DEH-trol el-ay)

## ABOUT OVERACTIVE BLADDER

Overactive bladder happens when the bladder muscle squeezes too often or it cannot be controlled. You may have wetting accidents (urge urinary incontinence). You may feel a strong need to pass urine right away (urgency). You may also feel you need to go often (frequency).

## WHO IS DETROL LA (long-acting) FOR?

Who can take DETROL LA?

Adults 18 years and older with symptoms of overactive bladder.

Who should not take DETROL LA?

Do not take DETROL LA if:

- You have trouble emptying your bladder (urinary retention).
- Your stomach empties slowly (gastric retention).
- You have an eye problem called uncontrolled narrow-angle glaucoma.
- You are allergic to anything in DETROL LA.

DETROL LA did not help the symptoms of overactive bladder when studied in children.

## BEFORE YOU START DETROL LA

Tell your doctor about all your medical conditions, including:

- Stomach or intestinal problems.
- Trouble emptying your bladder or if you have a weak urine stream.
- An eye problem called narrow-angle glaucoma.
- Liver or kidney problems.
- If you or any family members have a rare heart condition called QT prolongation (long QT syndrome). If you are pregnant or trying to become pregnant. We do not know if DETROL LA could harm your unborn baby.
- If you are breast-feeding. We do not know if DETROL LA passes into your milk or if it can harm your baby.

Tell your doctor about all your medicines. Include over-the-counter medicines, vitamins, and herbal products. Other drugs may change how your body handles DETROL LA. Your doctor may use a lower dose of DETROL LA if you take:

- Some medicines for fungus or yeast infections such as Nizoral<sup>®</sup> (ketoconazole), Sporanox<sup>®</sup> (itraconazole), and Monistat<sup>®</sup> (miconazole).
- Some medicines for bacteria infections such as Biaxin<sup>®</sup> (clarithromycin) and erythromycin.
- Sandimmune<sup>®</sup> (cyclosporine) or Velban<sup>®</sup> (vinblastine).

## POSSIBLE SIDE EFFECTS OF DETROL LA

The most common side effects are:

- dry mouth
- constipation
- headache
- stomach pain

These are not all the side effects of DETROL LA. For a complete list of side effects, ask your doctor or pharmacist.

## HOW TO TAKE DETROL LA

Do:

- Take DETROL LA exactly as your doctor tells you. For some people, DETROL LA may not work right away. Check with your doctor before stopping DETROL LA.
- Take DETROL LA capsules once a day with liquid, at the same time each day.
- Take DETROL LA with or without food.
- If you miss a dose of DETROL LA, begin taking DETROL LA again the next day. Do not take 2 doses of DETROL LA in the same day.
- Store DETROL LA at room temperature and out of light.

Don't:

- Do not drive a car or work with machines until you know how DETROL LA affects you. Medicines like DETROL LA can cause blurry vision. They can make you dizzy and sleepy.
- Do not change the dose on your own. Talk with your doctor.

## ABOUT DETROL LA

DETROL LA is a prescription medicine that treats the symptoms of overactive bladder.

- It reduces wetting accidents.
- It reduces the sudden, strong urge to pass urine.
- It helps you have fewer trips to the bathroom.

## NEED MORE INFORMATION?

- This is only a summary of important information. Ask your doctor or pharmacist for complete product information.
- Go to [www.detrolla.com](http://www.detrolla.com).
- Call (1-888) 4-DETROL (1-888-433-8765).



Distributed by:  
Pharmacia & Upjohn  
Division of Pfizer Inc, NY, NY 10017

Rx Only ©2006 Pfizer Inc All rights reserved. Printed in the USA.  
DEIF Rev 1 July 2006

Registered trademarks are the property of their respective owners.