

CHAPTER 9

Social mechanisms enhance escape responses in shoals of rainbowfish, *Melanotaenia duboulayi*

9.1 ABSTRACT

Shoaling behaviour has been shown to provide many benefits to group members including anti-predator and foraging advantages. In this study I examined the ability of fish shoals to escape from a novel trawl apparatus. Fish in shoals of 5 found, and escaped through, an escape hole in the oncoming trawl more quickly than fish in pairs. Fish in the larger shoals displayed a significant decrease in escape latencies over a series of five trawls, providing clear evidence of net avoidance learning, whereas fish in pairs showed no evidence of learning over successive runs. Observations suggested that more information on the location of the escape route was available to fish in larger shoal sizes. This occurred via social stimulation most probably gained through the observation other conspecifics in the shoal.

9.2 INTRODUCTION

Nets and other fishing gear invoke responses in fish resembling that of predator avoidance (Francis & Williams 1995) which has been shown to be facilitated by social learning (Suboski *et al.* 1990, Krause 1993, Mathis *et al.* 1996). Larger groups are thought to benefit from increased vigilance, which improves predator detection (Elgar 1989, Lima 1995). The phenomenon of collective detection assumes that all members of a group are alerted to the presence of a predator once a single group member has detected it (Lima 1995). Predator detection is followed by a fright response which rapidly spreads through the group, providing an early warning to those members that have not independently perceived the approaching threat (Godin & Morgan 1985, Godin *et al.* 1988). For example, in shoaling fish, fright responses move rapidly through the group resulting in synchronized schooling behaviours. The behaviour of an individual fish within a school is largely dependent on the behaviour of those around it. The speed of transmission of information through the school far exceeds that of the approaching predator (the 'Trafalgar effect': Treherne & Foster 1981). Shoaling fish thus provide an opportunity to study the flow of information between members of a group and how individuals monitor and respond to the behaviour of other group members.

A further benefit arising from the formation of groups is an increase in the probability that at least one member will discover a solution to a problem (e.g. show an effective escape response). Group members are more likely to make the "right" decision more quickly and with greater accuracy than they otherwise would in isolation (Grunbaum 1997). Increased shoal size facilitates information transfer and social learning which may be particularly important in detecting novel predators, exploiting novel food types or solving spatial problems. We might, therefore, hypothesise that larger groups of fish may solve novel problems sooner and the solution to the problem may spread more rapidly through the group. In this chapter I tested the hypotheses that by monitoring the behaviour of their shoal mates, fish in larger shoals: (1) discover the location of an escape route sooner, and (2) through successive exposure, learn the location of the escape route more rapidly.

9.3 **METHODS**

Female Duboulay's rainbowfish, *Melanotaenia duboulayi*, were captured using bait traps set in Amamoor Creek (26° 21'S, 152°40'E), a tributary of the Mary River in South-East Queensland. After being transported to the University of Queensland, the fish were held in a large holding tank (75cm x 60 cm x 25 cm deep) in the laboratory for several months prior to experimentation, allowing them ample time to adjust to captive conditions. After two days in captivity all fish readily accepted flake food. The average standard length of rainbowfish used during the experiment was 51.9 ± 1.4 mm (mean \pm SE).

An experimental tank was equipped with a pulley system that allowed a vertical net to be pulled along the long axis of the tank as in Chapter 2, Experiment 2 and Chapter 5. The depth of the water in the tank was maintained at 20 cm and the temperature at 23°C. The net had a mesh size of 1cm and completely blocked the tank with the exception of a small hole. The fish could use this hole to avoid being trapped as the net was dragged from one end to the other.

The apparatus and procedure are outlined in Chapter 2, Experiment 2 and Chapter 5. In this instance three people carried out the procedure. One observer called out the time as fish escaped through the hole and made general observations on the orientation and behaviour of the fish, another recorded the data, while a third wound the pulley handle. The first two experimenters sat motionless approximately two meters from the tank while the third was hidden from the fish behind both a barrier erected in the tank and a curtain.

Shoals of two and five fish were tested (10 replicates of each). The latency data were non-normally distributed because some fish did not escape from the trawl within the 90sec limit. A Wilcoxon Two-Sample test was used to examine the shoal size effect and a Kruskal-Wallis test (chi-square approximation) was used to examine the effect of run number for each shoal size. The statistics were performed using the SAS system (SAS Institute 1996).

9.4 **RESULTS**

Fish in groups of five consistently found the escape hole more quickly than fish in pairs (Figure 9.1). The number of exposures (run number) had a significant effect on escape latency for groups of five but not for groups of two over the five runs (Table 9.1, Figure 9.1). Groups of five fish decreased their mean (\pm SE) escape time from 74 ± 7 sec to 47 ± 5 seconds in five runs. By contrast, groups of two fish showed no significant learning effect and a larger amount of variability in escape latencies between runs.

During the first run all fish searched for an escape route. Of the 10 replicate groups, only one group of two successfully found the hole during the first run. By contrast, fish in groups of five discovered and escaped through the hole during their first exposure in 50% of the replicates. Once the net had stopped moving and was held in position, the intensity of searching increased until an individual succeeded in escaping. Although no quantitative information was collected on fish orientation, general observations suggested that the outcome of a trial appeared to vary depending on the orientation and location of the remaining fish. Fish that were orientated towards the escaping fish seemed to be attracted to the region where the hole was located. Fish that were not orientated towards escapees (and thus unable to monitor their behaviour) often remained trapped for the remainder of the run.




Figure 9.1. The mean escape latency in seconds (\pm SE) for fish in groups of two (triangles) and groups of five (squares) over five runs ($n=10$ in each case).

After four or five runs the groups of five swam with the net, keeping just in front of it. This behaviour is commonly seen in schools of fish avoiding trawls in the wild (Engas 1994). Close to the end of the tank they turned and swam calmly through the hole. Others simply moved to the far end of the tank, waited for the net to come to them, and then escaped. In contrast, the groups of two were more erratic. Rarely did a group of two find the hole by the end of the fourth or fifth run. Only 3 % of fish in pairs escaped the trawl while it was still in motion compared with 14.8 % of fish from groups of five. From our general observations it appeared that fish in the smaller groups spent most of their time with their heads facing away from the net as it moved. Fish in groups of five tended to be at varying distances and at different angles to the net, some facing it as it came towards them. Fish facing the net appeared to locate the hole sooner.

9.5 DISCUSSION

Rainbowfish in groups of five showed initial escape latencies that were significantly lower than fish in pairs. Furthermore, only the groups of five showed a rapid rate of learning with repeated exposure to the trawl. The only explanation for these results is that the higher levels of social interactions occurring in larger groups of fish facilitated more efficient escape responses.

Fish, like other vertebrates, show strong avoidance responses towards negative stimuli and quickly learn to avoid such stimuli in future encounters. Pyanov (1993), for example, showed that fish increase their defensive behaviour and learn to avoid trawls after just one exposure both in the laboratory and under natural conditions. Much of the learning that occurs in fish shoals may well be socially mediated and consequently dependent on shoal size. Hunter & Wisby (1964) investigated the net avoidance behaviour of carp using gear similar to that used in the present study and found that groups of fish were much more successful at avoiding capture than isolates.

(1) Shoal size effect:			
	S	Z	Prob > Z
Run 1	85.5	-1.771	0.077
Run 2	55.0	-3.998	<0.001
Run 3	75.0	-2.260	0.024
Run 4	56.0	-3.788	<0.001
Run 5	75.5	-2.452	0.014
Total	1658.0	-6.245	<0.001
(2) Run effect:			
	χ^2	Df	Prob > χ^2
Shoal size = 2	7.6128	4	0.107
Shoal size = 5	12.683	4	0.013

Table 9.1. Escape latency: (1) Results of Wilcoxon Two-Sample test (normal approximation, with continuity correction of 0.5) for shoal size effect between populations, and, (2) Kruskal-Wallis test (chi-square approximation) for run effect within each shoal size.

As with the carp in Hunter & Wisby's study, rainbowfish in groups of five often escaped while the net was still moving, whereas individuals were more likely to escape the net once it has stopped or failed to escape. Hunter & Wisby (1964) failed to pay close attention to the behaviour of individuals within the groups and accordingly could not explain the factors that may have led to the shorter reaction times observed over repeated runs in their study. From our observations it seems likely that the orientation of fish at the time of initial escape may be important in determining the fate of the rest of the group.

Hale (1956) found that even when "leadership" (i.e. the capacity of certain individuals to consistently perform better than others) was experimentally reduced, groups tended to escape more rapidly than singletons. Furthermore, the scores obtained when individuals were tested in groups bore no relationship to the scores obtained when those individuals were tested alone. These results indicate that improved group escape responses are the result of social interactions. In our study, the lower initial escape latency and faster improvement in performance observed in larger groups were unlikely to be the result of leaders or superior escapees.

Larger groups of fish tend to be bolder in the face of threat, and threat dilution appears to permit individuals to better concentrate on activities such as exploration and foraging (Magurran & Pitcher 1983, Smith & Warburton 1992). Welty (1934) suggested that goldfish escaped from a maze more quickly in a group due to elevated group cohesion, which resulted in increased exploration activity. Individuals in larger groups may have paid closer attention to the spatial details of the net and may have also detected the net coming towards them earlier, thus giving them more time to search for an escape route (early predator warning; Lazarus 1979).

Pitcher *et al.* (1982) showed that fish in larger shoals locate food faster than singletons. Information regarding the location of food is passed passively to shoal mates by a display of feeding behaviour, thus attracting conspecifics to the location (Pitcher & House 1987). The feeding responses of individuals are not independent, and social cues displayed by successful foragers inform other fish where to find food (Ryer & Olla 1992). The mechanisms underlying the responses observed in this experiment are probably very similar to those seen in forage patch copying. In this instance the cue for escape success may have been the movement of fish through

the net which attracted conspecifics to the general area of the escape hole. This social facilitation (Clayton 1978) probably contributed to the faster escape times for fish in larger shoals.

In the current experiment, the rapid decrease in escape latencies displayed by the fish in groups of five may have been influenced by individuals monitoring the behaviour of fellow shoal mates and benefiting from a discovery made by any one of them. This is supported by the fact that after the first fish escaped, others often followed in quick succession. Hale (1956) attributed group success in green sunfish (*Lepomis cyanellus*) to “mutual stimulation”. Fish grouping behaviour may also rely on guided social learning (Laland & Williams 1997) where the attention of an observer is attracted to a particular stimulus by a social model. The observer then interacts with the stimulus and relies on individual learning to complete the task. The rate of learning within a social group increases with the number of times the particular behaviour is successfully performed or the number of ‘tutors’ performing it (Giraldeau *et al.* 1994).

Fish that did not find the hole but followed others through it while escaping may have contributed less to the rediscovery of the escape route during subsequent runs. Previous experiments using groups of five fish indicated that the individual that located the hole first was more likely to escape during future runs (pers. obs). On many occasions members of smaller shoals found the hole during an early trial but failed to do so during the trials that followed it, perhaps because they had less assistance from other individuals in rediscovering it. More consistent searching responses are expected from larger shoals owing to the nature of the principal components of schooling behaviour (Grunbaum 1997). As shoal size increases inter-individual variation decreases due to behavioural averaging (Fitzsimmons & Warburton 1992). Such an effect may explain why the learning curve displayed by the fish in groups of five was not only steeper but also far smoother than that of fish in pairs.

Exploration and threat detection are facilitated by larger group sizes. Fish in larger shoals may also benefit from the discoveries made by their shoalmates. As shoal size increases the likelihood that an individual discovers the solution to a novel problem increases, as does the probability that another individual observes the solution being carried out. This results not only in a solution on first exposure to a problem, but also in an elevated rate of learning for the group as a whole.

[References](#)

[Back to Contents](#)