


Calculus I AP Workshop Problems
May 5, 2007

1. What is $\lim_{h \rightarrow 0} \frac{\cos\left(\frac{3\pi}{2} + h\right) - \cos\left(\frac{3\pi}{2}\right)}{h}$?

- (A) 1
- (B) $\frac{\sqrt{2}}{2}$
- (C) 0
- (D) -1
- (E) The limit does not exist.

2. At which of the five points on the graph in the figure at the right are $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$ both negative?



- (A) A
- (B) B
- (C) C
- (D) D
- (E) E

3. The slope of the tangent to the curve $y^3x + y^2x^2 = 6$ at $(2, 1)$ is

- (A) $-\frac{3}{2}$
- (B) -1
- (C) $-\frac{5}{14}$
- (D) $-\frac{3}{14}$
- (E) 0

4. A city is built around a circular lake that has a radius of 1 mile. The population density of the city is $f(r)$ people per square mile, where r is the distance from the center of the lake, in miles. Which of the following expressions gives the number of people who live within 1 mile of the lake?

- (A) $2\pi \int_0^1 r f(r) dr$
- (B) $2\pi \int_0^1 r (1 + f(r)) dr$
- (C) $2\pi \int_0^2 r (1 + f(r)) dr$
- (D) $2\pi \int_1^2 r f(r) dr$
- (E) $2\pi \int_1^2 r (1 + f(r)) dr$

Calculus I AP Workshop Problems
May 5, 2007

5. Which of the following statements about the function given by $f(x) = x^4 - 2x^3$ is true?
- (A) The function has no relative extremum.
 - (B) The graph of the function has one point of inflection and the function has two relative extrema.
 - (C) The graph of the function has two points of inflection and the function has one relative extremum.
 - (D) The graph of the function has two points of inflection and the function has two relative extrema.
 - (E) The graph of the function has two points of inflection and the function has three relative extrema.
6. If $f(x) = \sin^2(3 - x)$, then $f'(0) =$
- (A) $-2 \cos 3$
 - (B) $-2 \sin 3 \cos 3$
 - (C) $6 \cos 3$
 - (D) $2 \sin 3 \cos 3$
 - (E) $6 \sin 3 \cos 3$
7. The solution to the differential equation $\frac{dy}{dx} = \frac{x^3}{y^2}$, where $y(2) = 3$, is
- (A) $y = \sqrt[3]{\frac{3}{4}}x^4$
 - (B) $y = \sqrt[3]{\frac{3}{4}}x^4 + \sqrt[3]{15}$
 - (C) $y = \sqrt[3]{\frac{3}{4}}x^4 + 15$
 - (D) $y = \sqrt[3]{\frac{3}{4}}x^4 + 5$
 - (E) $y = \sqrt[3]{\frac{3}{4}}x^4 + 15$
8. What is the average rate of change of the function f given by $f(x) = x^4 - 5x$ on the closed interval $[0, 3]$?
- (A) 8.5
 - (B) 8.7
 - (C) 22
 - (D) 33
 - (E) 66
9. The position of a particle moving along a line is given by $s(t) = 2t^3 - 24t^2 + 90t + 7$ for $t \geq 0$. For what values of t is the speed of the particle increasing?
- (A) $3 < t < 4$ only
 - (B) $t > 4$ only
 - (C) $t > 5$ only
 - (D) $0 < t < 3$ and $t > 5$
 - (E) $3 < t < 4$ and $t > 5$

Calculus I AP Workshop Problems
May 5, 2007

10. $\int (x - 1)\sqrt{x} \, dx =$

(A) $\frac{3}{2}\sqrt{x} - \frac{1}{\sqrt{x}} + C$

(B) $\frac{2}{3}x^{\frac{3}{2}} + \frac{1}{2}x^{\frac{1}{2}} + C$

(C) $\frac{1}{2}x^2 - x + C$

(D) $\frac{2}{5}x^{\frac{5}{2}} - \frac{2}{3}x^{\frac{3}{2}} + C$

(E) $\frac{1}{2}x^2 + 2x^{\frac{3}{2}} - x + C$

11. What is $\lim_{x \rightarrow \infty} \frac{x^2 - 4}{2 + x - 4x^2}$?


(A) -2

(B) $-\frac{1}{4}$

(C) $\frac{1}{2}$

(D) 1

(E) The limit does not exist.



12. The figure above shows the graph of $y = 5x - x^2$ and the graph of the line $y = 2x$. What is the area of the shaded region?

(A) $\frac{25}{6}$

(B) $\frac{9}{2}$

(C) 9

(D) $\frac{27}{2}$

(E) $\frac{45}{2}$

Calculus I AP Workshop Problems
 May 5, 2007

13. If f is a function that is continuous for all real numbers, then


$$\frac{d}{dx} \int_0^{x^2} f(t) dt =$$

- (A) $2x f(x^2)$
- (B) $2x f(2x)$
- (C) $f(2x)$
- (D) $f(x^2)$
- (E) $f'(x^2)$


14. Which of the following is a slope field for the differential equation

$$\frac{dy}{dx} = \frac{x}{y}?$$


(A)




(B)




(C)



(D)



(E)



Calculus I AP Workshop Problems
May 5, 2007

15. A particle travels along a straight line with a velocity of $v(t) = 3e^{(-t/2)} \sin(2t)$ meters per second. What is the total distance, in meters, traveled by the particle during the time interval $0 \leq t \leq 2$ seconds?

- (A) 0.835
- (B) 1.850
- (C) 2.055
- (D) 2.261
- (E) 7.025

16. Let S be the region enclosed by the graphs of $y = 2x$ and $y = 2x^2$ for $0 \leq x \leq 1$. What is the volume of the solid generated when S is revolved about the line $y = 3$?

(A) $\pi \int_0^1 \left((3 - 2x^2)^2 - (3 - 2x)^2 \right) dx$


(B) $\pi \int_0^1 \left((3 - 2x)^2 - (3 - 2x^2)^2 \right) dx$

(C) $\pi \int_0^1 (4x^2 - 4x^4) dx$

(D) $\pi \int_0^2 \left(\left(3 - \frac{y}{2} \right)^2 - \left(3 - \sqrt{\frac{y}{2}} \right)^2 \right) dy$

(E) $\pi \int_0^2 \left(\left(3 - \sqrt{\frac{y}{2}} \right)^2 - \left(3 - \frac{y}{2} \right)^2 \right) dy$

17.



The graph of a function f is shown above. If $\lim_{x \rightarrow b} f(x)$ exists and f is not continuous at b , then $b =$

- (A) -1
- (B) 0
- (C) 1
- (D) 2
- (E) 3

Calculus I AP Workshop Problems
May 5, 2007

18.

x	1.1	1.2	1.3	1.4
$f(x)$	4.18	4.38	4.56	4.73

Let f be a function such that $f''(x) < 0$ for all x in the closed interval $[1, 2]$. Selected values of f are shown in the table above. Which of the following must be true about $f'(1.2)$?


- (A) $f'(1.2) < 0$
- (B) $0 < f'(1.2) < 1.6$
- (C) $1.6 < f'(1.2) < 1.8$
- (D) $1.8 < f'(1.2) < 2.0$
- (E) $f'(1.2) > 2.0$

19.

Two particles start at the origin and move along the x -axis. For $0 \leq t \leq 10$, their respective position functions are given by $x_1 = \sin t$ and $x_2 = e^{-2t} - 1$. For how many values of t do the particles have the same velocity?

- (A) None
- (B) One
- (C) Two
- (D) Three
- (E) Four

20.



Graph of f






The graph of the function f shown above consists of two line segments. If g is the function defined by $g(x) = \int_0^x f(t) dt$, then

$g(-1) =$

- (A) -2
- (B) -1
- (C) 0
- (D) 1
- (E) 2

Calculus I AP Workshop Problems
May 5, 2007

21. The graphs of five functions are shown below. Which function has a nonzero average value over the closed interval $[-\pi, \pi]$?



22. The base of a solid S is the semicircular region enclosed by the graph of $y = \sqrt{4 - x^2}$ and the x -axis. If the cross sections of S perpendicular to the x -axis are squares, then the volume of S is

- (A) $\frac{32\pi}{3}$
 (B) $\frac{16\pi}{3}$
 (C) $\frac{40}{3}$
 (D) $\frac{32}{3}$
 (E) $\frac{16}{3}$

Calculus I AP Workshop Problems
May 5, 2007

23. Oil is leaking from a tanker at the rate of $R(t) = 2,000e^{-0.2t}$ gallons per hour, where t is measured in hours. How much oil leaks out of the tanker from time $t = 0$ to $t = 10$?
- (A) 54 gallons
(B) 271 gallons
(C) 865 gallons
(D) 8,647 gallons
(E) 14,778 gallons
24. If $f'(x) = \sin\left(\frac{\pi e^x}{2}\right)$ and $f(0) = 1$, then $f(2) =$
- (A) -1.819
(B) -0.843
(C) -0.819
(D) 0.157
(E) 1.157

Answers to Calculus AB Multiple-Choice Questions

Part A

- | | | |
|------|-------|-------|
| 1. A | 6. B | 11. B |
| 2. B | 7. E | 12. B |
| 3. C | 8. C | 13. A |
| 4. D | 9. E | 14. E |
| 5. C | 10. D | |

Part B

- | | | |
|--------|--------|--------|
| 15.* D | 19.* D | 22. D |
| 16. A | 20. B | 23.* D |
| 17. B | 21. E | 24.* E |
| 18. D | | |

* Indicates a graphing calculator-active question.