

LIVING ANIONIC POLYMERIZATION OF 2-  
VINYLNAPHTHALENE. SYNTHESIS AND CHARACTERIZATION  
OF MACROCYCLIC POLY(2-VINYLNAPHTHALENE)

by

Gennadi Nossarev

---

A Dissertation Presented to the  
FACULTY OF THE GRADUATE SCHOOL  
UNIVERSITY OF SOUTHERN CALIFORNIA  
In Partial Fulfillment of the  
Requirements for the Degree  
DOCTOR OF PHILOSOPHY  
(CHEMISTRY)

December 2001

Copyright 2001

Gennadi Nossarev

## Table of Contents

Dedication	ii
Acknowledgements	iii
List of Tables	viii
List of Schemes	ix
List of Figures	x
<b>Chapter 1. General Introduction</b>	
1.1 Light Harvesting in Natural Systems	1
1.2 Experimental Modeling of Light Harvesting Processes	3
1.3 Macrocyclic Vinylaromatic Polymers as Promising Light Harvesting Models	4
1.4 Synthetic Methodologies for Preparation of Macrocyclic Polymers	6
1.5 References	10
<b>Chapter 2. Experimental Section</b>	
2.1 High Vacuum Line Techniques	19
2.2 Purification and Treatment of Reagents	20
2.3 Preparation of Initiators	28
2.4 Polymerization, Block-Copolymerization, End Capping and Coupling Reactions under High Vacuum	30
2.5 Polymerization, End Capping and Coupling Reactions under Argon	32
2.6 Polymerization and Cyclization Reactions under High Vacuum	33
2.7 Polymer Characterization and Analysis	35
2.7.1 Size Exclusion Chromatography	35

2.7.2	Nuclear Magnetic Resonance	36
2.7.3	UV-Vis Absorption and Fluorescence Spectroscopy	37
2.7.4	Differential Scanning Calorimetry	38
2.7.5	Thermal Gravimetric Analysis	38
2.7.6	Evidence for Macrocyclic Polymers	38
2.7	References	40

### Chapter 3. Living Anionic Polymerization of 2-Vinylnaphthalene in Tetrahydrofuran

3.1	Introduction	48
3.2	Experimental	50
3.3	Polymerizations in the Presence of Lithium Ion. Role of Monomer Purification	51
3.4	Potassium Ion Mediated Living Anionic Polymerization of 2-Vinylnaphthalene	54
3.4.1	Formation of Block Copolymers	57
3.4.2	End- functionalization Reactions	60
3.5	References	61

### Chapter 4. Living t-BuLi Initiated Anionic Polymerization of 2-Vinylnaphthalene in Toluene-Tetrahydrofuran Mixtures

4.1	Introduction	77
4.2	Experimental	78
4.3	Kinetic Studies of 2VN Polymerization	79
4.4	End-functionalization and Coupling Reactions	83

4.5	Molecular Weight Determination	84
4.6	References	86
<b>Chapter 5. Synthesis and Characterization of Macrocyclic Poly (2-vinylnaphthalene) Containing Single 1,4-Benzylidene or 9,10-Anthracenylidene Groups</b>		
5.1	Introduction	101
5.2	Experimental	102
5.3	Macrocyclic P2VN Containing Single 1,4-Benzylidene Linkage	103
5.4	Macrocyclic P2VN Containing Single 9,10-Anthracenylidene Linkage	108
5.5	Spectroscopic Studies of Macrocyclic and Linear P2VN	110
5.6	P2VN as Hole Transporting Materials in OLED's	115
5.7	References	119
	Bibliography	142
	Appendix	149

## List of Tables

<b>Table 3.1</b>	Effect of the monomer impurity concentrations on P2VN-Li end-functionalization.	64
<b>Table 3.2</b>	Resumption polymerizations in THF at $-78\text{ }^{\circ}\text{C}$ with purified 2VN.	65
<b>Table 3.3</b>	Radical anion initiated polymerizations of 2-vinylnaphthalene in THF at $-78\text{ }^{\circ}\text{C}$ .	66
<b>Table 3.4</b>	Restart and block-copolymerization involving P2VN- $\text{K}_2$ .	67
<b>Table 3.5</b>	Coupling reactions of P2VN- $\text{K}_2$ and DBX in THF at $-78\text{ }^{\circ}\text{C}$ .	68
<b>Table 4.1</b>	Polymerization of 2VN in toluene/THF. Effect of THF concentration on the polymerization kinetics.	88
<b>Table 4.2</b>	Polymerization of 2VN in toluene/THF. End-functionalization and coupling reactions of P2VN-Li	89
<b>Table 4.3</b>	Comparison of P2VN molecular weight determined by SEC and $^1\text{H}$ NMR.	90
<b>Table 5.1</b>	Formation of macrocyclic P2VN by coupling of P2VN- $\text{K}_2$ and DBX or BCMA in THF at $-78\text{ }^{\circ}\text{C}$ .	122
<b>Table 5.2</b>	Thermal properties of fractionated macrocyclic and matching linear P2VN containing single 1,4 benzylidene or 9,10-anthracenylidene units.	123
<b>Table 5.3</b>	Polymers used as hole transporting materials and characteristics of devices prepared with these polymers.	124

## List of Schemes

<b>Scheme 1.1</b>	Synthesis of macrocyclic vinylaromatic polymers with variations of chromophores at initiator site, in repeating unit and in coupling group.	12
-------------------	---	----

## List of Figures

<b>Figure 1.1</b>	Energy transfer upon quenching of a chromophore excited state.	13
<b>Figure 1.2</b>	Structure of light-harvesting system in photosynthetic bacteria and the path for energy transfer from LH1 to a special pair of bacteriochlorophylls in RC.	14
<b>Figure 1.3</b>	An example of covalently linked chromophore-quencher assembly.	15
<b>Figure 1.4</b>	Covalently bound polymer system of donor ( <b>D</b> ) and acceptor ( <b>A</b> ).	16
<b>Figure 1.5</b>	Simplified representation of the synthesis of macrocyclic polymers by anionic polymerization and end-to-end coupling of polymer dianions with bifunctional electrophiles.	17
<b>Figure 1.6</b>	Sybyl calculated minimized structure of isotactic poly(2-vinylnaphthalene) with $DP_n = 10$ .	18
<b>Figure 2.1</b>	Typical set up for distillation of toluene or similar low boiling point liquids under high vacuum.	41
<b>Figure 2.2</b>	Apparatus for formation of a mirror of an alkali metal under high vacuum for initiator preparation or thorough solvent purification.	42
<b>Figure 2.3</b>	Set up for pre-purification of 2-vinylnaphthalene by sublimation from a mild scavenger ( $CaH_2$ ) under high vacuum.	43
<b>Figure 2.4</b>	Set up for rigorous purification of 2-vinylnaphthalene by sublimation from a scavenger ( $CaH_2$ or $LiAlH_4$ ) under high vacuum.	44
<b>Figure 2.5</b>	Typical apparatus for polymerization and consequent reactions involving polymer carbanion under high vacuum.	45

<b>Figure 2.6</b>	Typical apparatus for polymerizations under ultra pure inert gas.	46
<b>Figure 2.7</b>	Apparatus for polymerization and consequent cyclization of polymer dianion under high vacuum.	47
<b>Figure 3.1</b>	SEC curves of poly(2-vinylnaphthalene) prepared using sublimed <i>t</i> -BuLi and monomer purified a) by 2 sublimations from CaH <sub>2</sub> (Table 3.1, exp. # 8) and b) by sublimation from LiAlH <sub>4</sub> followed by sublimation from CaH <sub>2</sub> (Table 3.2, exp. # 97).	69
<b>Figure 3.2</b>	250 MHz <sup>1</sup> H NMR of 2-vinylnaphthalene in CDCl <sub>3</sub> a) sublimed 2 times from CaH <sub>2</sub> and b) sublimed from LiAlH <sub>4</sub> followed by sublimation from CaH <sub>2</sub> .	70
<b>Figure 3.3</b>	62.9 MHz <sup>13</sup> C NMR of 2-vinylnaphthalene in CDCl <sub>3</sub> sublimed 2 times from CaH <sub>2</sub> : a) the whole spectrum, b) the region showing carbonyl and c) methyl resonances.	71
<b>Figure 3.4</b>	SEC curves of poly(2-vinylnaphthalene) from exp. # 88 (Table 3.2): a) P2VN precursor and b) P2VN after addition of the second 2VN portion after 1 h of storage at -78 °C (peak at 19 mL corresponds to OMOHN).	72
<b>Figure 3.5</b>	Normalized SEC curves of P2VN (Table 3.4, # 87): a) P2VN precursor and b) P2VN after addition of the second 2VN portion after 1 h of storage (peak at 19 mL corresponds to OMOHN).	73
<b>Figure 3.6</b>	Normalized SEC curves of P2VN and PDMVF- <i>b</i> -P2VN- <i>b</i> -PDMVF (Table 3.4, # 91) (peak at 19 mL corresponds to OMOHN).	74
<b>Figure 3.7</b>	250 MHz <sup>1</sup> H NMR of a) P2VN (Table 3.4, # 94), b) PDMVF- <i>b</i> -P2VN- <i>b</i> -PDMVF (Table 3.4, # 91) and c) PAMS- <i>b</i> -P2VN- <i>b</i> -PAMS (Table 3.4, # 94).	75
<b>Figure 3.8</b>	Normalized SEC curves of a) P2VN and P2VN- <i>b</i> -PAMS- <i>b</i> -P2VN (Table 3.4, # 95), b) P2VN and PAMS- <i>b</i> -P2VN- <i>b</i> -PAMS (Table 3.4, # 94).	76

<b>Figure 4.1</b>	Normalized SEC chromatograms of P2VN initiated with <i>t</i> -BuLi in toluene at room temperature (Table 4.1, # 57) and terminated with MeOH after: a) 30 min, b) 60 min, c) 120 min, d) 180 min.	91
<b>Figure 4.2</b>	Normalized SEC chromatograms of P2VN initiated with <i>t</i> -BuLi in toluene in the presence of 31 mM THF at $-78\text{ }^{\circ}\text{C}$ (Table 4.1, # 55) and terminated with MeOH after: a) 5 min, b) 15 min, c) 45 min, d) 180 min.	92
<b>Figure 4.3</b>	Normalized SEC chromatograms of P2VN initiated with <i>t</i> -BuLi in toluene in the presence of 58 mM THF at $-78\text{ }^{\circ}\text{C}$ (Table 4.1, # 56) and terminated with MeOH after: a) 5 min, b) 15 min, c) 45 min, d) 180 min.	93
<b>Figure 4.4</b>	<i>t</i> -BuLi initiated polymerization of 2VN in toluene/THF. Dependence of $\text{Ln}[1-(I_0/M_0)\times\text{DP}_n]$ vs. time: a) Table 4.1, # 55, b) Table 4.1, # 105 and c) Table 4.1, # 56.	94
<b>Figure 4.5</b>	Relationship between number-average MW and monomer conversion for 2VN polymerization in toluene/THF (Table 4.1, # 105).	95
<b>Figure 4.6</b>	Normalized SEC chromatograms of P2VN initiated with <i>t</i> -BuLi in toluene/THF and coupled with DBX: a) Table 4.2, # 65 and b) Table 4.2, # 66.	96
<b>Figure 4.7</b>	250 MHz proton NMR spectrum in $\text{CDCl}_3$ of P2VN-Li terminated with TMSCl (Table 4.2, # 63).	97
<b>Figure 4.8</b>	UV-Vis spectrum acquired in 10 mm quartz cell in cyclohexane/(20% v.) THF of a) P2VN terminated with MeOH ( $\text{DP}_n = 60$ , $D = 1.12$ , conc. = 40 mg/L), b) P2VN coupled with BCMA (Table 4.2, # 62, $\text{DP}_n = 67$ , $D = 1.13$ , conc. = 48 mg/L) and c) the trace b magnified 10 times.	98

<b>Figure 4.9</b>	Emission spectrum acquired in 10 mm quartz cell in cyclohexane/(20% v.) THF of a) P2VN terminated with MeOH ( $DP_n = 60$ , $D = 1.12$ , conc. = 48 mg/L), b) P2VN coupled with BCMA (Table 4.2, # 62, $DP_n = 67$ , $D = 1.13$ , conc. = 48 mg/L) and c) the spectrum magnified 5 times.	99
<b>Figure 4.10</b>	250 MHz proton NMR spectrum in $CDCl_3$ of P2VN-Li coupled with DCDMS (Table 4.2, # 74).	100
<b>Figure 5.1</b>	Normalized SEC (RI) traces of a) precursor P2VN and b) reaction product of high anion concentration coupling with DBX (Table 5.1, # 80).	125
<b>Figure 5.2</b>	Normalized SEC (RI) traces of a) precursor P2VN, b) product of coupling with DBX under high dilution and c) fractionated macrocyclic P2VN (Table 5.1, # 92).	126
<b>Figure 5.3</b>	Relationship between glass transition temperature ( $T_g$ ) and degree of polymerization of a) linear and b) macrocyclic P2VN (the data point of linear P2VN of a $DP_n = 10$ (open diamond) is taken from reference 18).	127
<b>Figure 5.4</b>	Differences in thermal non-oxidative decomposition (TGA) for linear(solid line) and macrocyclic P2VN (dashed line) with $DP_n$ values of a) 18 (Table 5.2, # 83) and b) 125 (Table 5.2, # 85).	128
<b>Figure 5.5</b>	Normalized SEC (RI) traces of a) linear precursor and b) unfractionated product of high dilution coupling of P2VN- $K_2$ with BCMA (Table 5.1, # 102).	129
<b>Figure 5.6</b>	Normalized SEC traces of a) precipitated linear and b) fractionated macrocyclic P2VN acquired using RI and c) UV at 405 nm detectors (Table 5.2, # 102).	130
<b>Figure 5.7</b>	Absorption spectrum of fractionated macrocyclic P2VN with $DP_n=39$ (Table 5.2, # 102). containing 9,10-anthracenylidene group (54 mg/L in cyclohexane/THF : 80/20, v/v), 0.1 cm cell). Insert: is magnified 10 $\times$ .	131

<b>Figure 5.8</b>	Absorption spectra of a) 2-vinylnaphthalene (2VN) sublimed once, b) 2-ethylnaphthalene (2EN) (50 mg/L in cyclohexane/THF = 90/10 v/v, 0.1 cm cell) and c) magnified region of S <sub>1</sub> transition.	132
<b>Figure 5.9</b>	Emission spectra of a) 2-ethylnaphthalene (2EN) with excitation at 280 nm and b) excitation at 290 nm, c) 2-vinylnaphthalene sublimed once with excitation at 280 nm and d) excitation at 290 nm (50 mg/L in cyclohexane/THF = 90/10 v/v, 1 cm cell, slit 3.0 nm).	133
<b>Figure 5.10</b>	Absorption spectra of a) linear P2VN (50 mg/L) and b) macrocyclic P2VN (53 mg/L) with DP <sub>n</sub> =12 (Table 5.2, # 99) and c) 2EN (50 mg/L) in cyclohexane/THF = 90/10 v/v, 1 cm cell.	134
<b>Figure 5.11</b>	Emission spectra of a) linear P2VN (26 mg/L) and b) macrocyclic P2VN (26 mg/L) with DP <sub>n</sub> =18 in cyclohexane/THF =95/5 v/v, 1 cm cell, slit 3.0 nm, excitation at 280 nm (Table 5.2, # 83).	135
<b>Figure 5.12</b>	Emission spectra of polymers with DP <sub>n</sub> =7 a) linear P2VN, b) macrocyclic P2VN and c) macrocyclic P2VN after hydrogenation, 50 mg/L in cyclohexane/THF = 90/10 v/v, 1 cm cell, slit 6.0 nm, excitation at 285 nm (Table 5.1, # 109).	136
<b>Figure 5.13</b>	Emission spectra of a) linear P2VN (50 mg/L) and b) macrocyclic P2VN (53 mg/L) with DP <sub>n</sub> =12 in cyclohexane/THF = 90/10 v/v, 1 cm cell, slit 3.0 nm, excitation at 285 nm (Table 5.2, # 99).	137
<b>Figure 5.14</b>	Emission spectra of a) linear P2VN, b) macrocyclic P2VN with DP <sub>n</sub> =17 and c) 2EN, 54 mg/L in cyclohexane/THF = 90/10 v/v/, 1 cm cell, slit 3.0 nm, excitation at 280 nm (Table 5.2, # 92).	138
<b>Figure 5.15</b>	Monomer-to-excimer band ratios in emission spectra as a function of DP <sub>n</sub> for a) linear P2VN and b) macrocyclic P2VN in cyclohexane/THF =90/10 v/v, 1 cm cell, slit 3.0 nm, excitation at 285 nm (Table 5.2).	139

<b>Figure 5.16</b>	Quantum efficiency of OLED's with hole transporting layer spin coated using polymers in Table 5.3: a) # 1, b) # 4, c) # 7, d) # 3 and e) # 8.	140
<b>Figure 5.17</b>	Current – voltage characteristics of OLED's with hole transporting layers spin coated using polymers in Table 5.3: a) # 8, b) # 3, c) # 7, d) # 4 and e) #1.	141