

NUMERICAL SOLUTIONS OF THE ALGEBRAIC MATRIX RICCATI EQUATION

HANS M. AMMAN AND HEINZ NEUDECKER

Date 2-9-1997

ABSTRACT.

The Linear-Quadratic control model is one of the most widely used control models in both empirical and theoretical economic modeling. In order to obtain the equilibrium solution of this control model, the so-called algebraic matrix Riccati equation, has to be solved. In this note we present a numerical solution method for solving this equation. Our method solves the Riccati equation as a multidimensional fixed-point problem. By establishing the analytical derivative of the Riccati equation we have been able to construct a very efficient Newton-type solution method with quadratic convergence properties. Our method is an extension for the Newton-Raphson method described in Kwakernaak and Sivan (1972) and does not require any special conditions on the transition matrix as in the nonrecursive method of Vaughan (1970).

INTRODUCTION

One of the most widely used control models in economics is the Linear-Quadratic control model. Recent examples are found in McGratten (1994), Amman and Kendrick (1995) and many more. Generally, the solution of this control model for finite-horizon problems is obtained by solving the so-called Riccati matrix equation backward in time from the terminal date to an initial date. Solving the Riccati equation recursively in time is a simple operation which generally does not pose any difficulties. The procedure is different, however, for infinite-horizon problems. In that case the algebraic matrix Riccati equation has to be solved. Unfortunately, there seems to be no analytical solution to this equation.

Bertsekas (1976) pointed out that under certain restrictions the algebraic matrix Riccati equation could be solved by using successive substitutions and that it has a unique solution. Successive substitutions are quite inefficient, however, and it may take many iteration steps before a solution is reached. In this note we will present a fast method for solving the algebraic Riccati equation numerically, using a Newton-type solution method. The advantage of such a method, as generally known, is that it has quadratic convergence properties ensuring a rapid solution.

Key words and phrases. Algebraic Matrix Riccati equation, Newton Numerical solution method, Equilibrium Solution. *JEL code* C61, C63.

Our special thanks go to Ken Judd and Berc Rustem, who gave us valuable advice on an earlier version of the paper. The corresponding author is Hans M. Amman, Department of Economics, University of Amsterdam, Roetersstraat 11, 1018 WB Amsterdam, the Netherlands. Email amman@sara.nl.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ - \TeX

Our method deviates from the newer approaches in the engineering literature, see Lainiotis et al. (1994) and Stoorvogel and Weeren (1994)¹. These algorithms are of a recursive nature and do not exploit derivative information.

A nonrecursive method for solving the algebraic Riccati equation, popular in economics, is the method of Vaughan (1970). Unfortunately this method is only applicable to a limited set of problems as it requires that the transition matrix should be nonsingular. For most applications in economics this conditions does not hold and an alternative method should be used. In the next paragraph we present such a method that does not require any additional assumptions on the matrices in the Linear-Quadratic control model. Our method is an extension of the Newton-Raphson method described in Kwakernaak and Sivan (1972) in the sense that we use derivative information for solving the Riccati equation. Also, it does not require any special conditions on the transition matrix as in the nonrecursive method of Vaughan (1970).

THE RICCATI EQUATION

Following Kendrick (1981), a commonly used form of the Linear-Quadratic control model is to minimize

$$J = \frac{1}{2}[x_T - \hat{x}_T]'W_T[x_T - \hat{x}_T] + \frac{1}{2}\sum_{t=0}^{T-1}\{[x_t - \hat{x}_t]'W_t[x_t - \hat{x}_t] + [u_t - \hat{u}_t]'R_t[u_t - \hat{u}_t] + [x_t - \hat{x}_t]'F_t[u_t - \hat{u}_t]\} \quad (1)$$

subject to the constraints

$$x_{t+1} = A_t x_t + B_t u_t + c_t \quad (2)$$

x_0 given

with $t \in \{0, 1, \dots, T\}$, $x_t \in \mathfrak{R}^n$ as the state vector, $u_t \in \mathfrak{R}^m$ as the control vector, $c_t \in \mathfrak{R}^n$ the vector of exogenous variables, $\hat{x}_t \in \mathfrak{R}^n$ and $\hat{u}_t \in \mathfrak{R}^m$ as target vectors, $A_t \in \mathfrak{R}^{(n \times n)}$ and $B_t \in \mathfrak{R}^{(n \times m)}$ as system matrices and $W_t \in \mathfrak{R}^{(n \times n)}$, $R_t \in \mathfrak{R}^{(m \times m)}$, $F_t \in \mathfrak{R}^{(n \times m)}$ as penalty matrices. Normally W_t and R_t are assumed to be positive semi-definite and positive definite respectively. In this general form of the Linear-Quadratic control model, the Riccati equation has the form

$$X_t = W_{t+1} + A_t' X_{t+1} A_t - (A_t' X_{t+1} B_t + F_t)(R_t + B_t' X_{t+1} B_t)^{-1} (B_t' X_{t+1} A_t + F_t') \quad (3)$$

with the fixed end condition

$$X_T = W_T \quad (4)$$

¹We would like to thank one of the anonymous referees for bringing to our attention a number of recent references in this area.

Following Sargent (1987) or Lancaster and Rodman (1995), the infinite-horizon equilibrium solution of the Riccati equation, if the matrices are time-invariant, leads to the algebraic matrix Riccati equation

$$X = W + A'XA - (A'XB + F)(R + B'XB)^{-1}(B'XA + F') \quad (5)$$

Let us now define the following function

$$G(X) = X - W - A'XA + (A'XB + F)(R + B'XB)^{-1}(B'XA + F') \quad (6)$$

This leads to the first derivative

$$\begin{aligned} \frac{\partial g}{\partial z'} &= D^+[I - \{(A'XB + F)(R + B'XB)^{-1}B' - A'\} \otimes \\ &\quad \{(A'XB + F)(R + B'XB)^{-1}B' - A'\}]D \end{aligned} \quad (7)$$

with $g = D^+ \text{vec } G$ and $z = D^+ \text{vec } X$. The second derivatives have the form

$$\begin{aligned} H_k &= \frac{\partial^2 g_k}{\partial z \partial z'} = D'[A\tilde{E}_{ij}A' \otimes B(R + B'XB)^{-1}B']D \\ &\quad - D'[A\tilde{E}_{ij}(A'XB + F)(R + B'XB)^{-1}B' \otimes B(R + B'XB)^{-1}B']D \\ &\quad - D'[B(R + B'XB)^{-1}(B'XA + F')\tilde{E}_{ij}A' \otimes B(R + B'XB)^{-1}B']D \\ &\quad + D'[B(R + B'XB)^{-1}(B'XA + F')\tilde{E}_{ij}(A'XB + F) \\ &\quad \quad (R + B'XB)^{-1}B' \otimes B(R + B'XB)^{-1}B']D \end{aligned} \quad (8)$$

where the matrix I is the identity matrix and \otimes denotes the Kronecker product and $\tilde{E}_{ij} = E_{ij} + E_{ji}$, $i, j \in \{1, \dots, n\}$. The matrices D and D^+ are the duplication matrix and inverse duplication matrix (cf. Magnus and Neudecker (1994)). These matrices "remove and restore" redundant upper triangular parts of the (symmetric) positive definite Riccati matrix X and the function $G(X)$. As a result the matrix X is transformed to a $p = \frac{1}{2}n(n+1)$ vector z . Consequently, H_k is a $(p \times p)$ matrix of second derivatives of the k th element g_k of the vector $g(z)$ with respect to the vector z , k being $(j-1)n + i$.

The problem here is to solve the multi-dimensional fixed-point problem $g(z) = 0$. There are several types of numerical methods to solve such a problem. With the help of equation (7) it is simple to construct an iterative scheme to solve our fixed-point problem, *viz*

$$z^{(j+1)} = z^{(j)} - \left(\frac{\partial g}{\partial z'}\right)^{-1}g \quad (9)$$

with $z^{(0)} = D^+ \text{vec } W$ as an initial condition. This updating scheme is quite robust and will generally lead to a rapid solution up to some machine precision level. However, this method will in some particular cases not converge. This can be shown by transforming our problem to a nonlinear least-squares problem of the form (see Press et al. (1986)),

$$\min : \psi(z) = \frac{1}{2}g(z)'g(z) \quad (10)$$

which has the first derivative

$$\frac{\partial \psi}{\partial z'} = g' \left(\frac{\partial g}{\partial z'} \right) \quad (11)$$

and the second derivative

$$\frac{\partial^2 \psi}{\partial z \partial z'} = \left(\frac{\partial g}{\partial z'} \right)' \left(\frac{\partial g}{\partial z'} \right) + \sum_{j=1}^n \sum_{i=j}^n g_k H_k, \quad k = (j-1)n + i \quad (12)$$

which is the Hessian matrix of $\psi(z)$ with respect to z , k , g_k and H_k as previously defined. Clearly, the Hessian matrix is positive (semi) definite in the neighbourhood of the optimal solution $g(z) = 0$. As such equation (12) serves as a diagnostic on convergence of our multidimensional fixed-point algorithm. Furthermore, if the initial value of $g(z^{(0)})$ deviates substantially from zero, nonconvergence may occur. This convergence issue can be fixed to a large extent by adding a relaxation factor λ to equation (9), which leads to

$$z^{(j+1)} = z^{(j)} - \lambda \left(\frac{\partial g}{\partial z'} \right)^{-1} g \quad (13)$$

The relaxation factor λ is chosen such that equation (10) is globally optimized. There are several strategies described in Press et al. (1986) for choosing λ efficiently.

TWO EXAMPLES

Let's assume we have the following matrices that belong to a simple control model with $n = 5$ and $m = 1$, see Amman (1995)²,

$$A = \begin{pmatrix} 0.8 & 0 & 0 & 0 & -0.8 \\ 0.8 & 0 & 0 & 0 & -0.8 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0.25 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$B = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \\ 0 \end{pmatrix},$$

$F = 0$, $W = I$ and $R = 1$ which leads to the solution of equation (5)

$$X = \begin{pmatrix} 2.2069 & 0 & 0 & 0 & -1.1976 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ -1.1976 & 0 & 0 & 0 & 2.3115 \end{pmatrix}$$

²A GAUSS implementation of the algorithms can be obtained through email from the corresponding author.

Clearly, this example could not be solved by using Vaughan's (1970) nonrecursive method due to the fact that the A matrix is singular.

Table 1³ below gives the norm $\|g(z^{(j)})\| = \sqrt{g(z^{(j)})'g(z^{(j)})}$ at each iteration step for the above model when we use the Newton method of equation (9). As expected the Newton method converges in a small number of iterations.

TABLE 1

j	$\ g(z^{(j)})\ $
1	1.1921E-01
2	2.7930E-05
3	5.3938E-13
4	7.6919E-16
∞	2.2404E-16

However, if the matrix A has the form⁴

$$A = \begin{pmatrix} -0.5208 & 0.5999 & -0.4380 & -0.3014 & -0.0562 \\ 0.9405 & -1.7373 & -1.5401 & -2.1367 & -1.2417 \\ 0.1110 & -0.8929 & -0.5187 & -1.8992 & -2.1634 \\ 0.0058 & 0.9553 & -0.8661 & -0.7301 & 1.7773 \\ 1.0474 & 0.5714 & 0.3946 & 1.1376 & -0.4130 \end{pmatrix},$$

the Newton method of equation (9) will not converge. The eigenvalues of the Hessian matrix in (12) for the initial value $X = W$ are in descending order $\{430.343, 104.7753, 36.3720, 15.4705, 8.8303, 7.8615, 4.7562, 1.3750, 1.1323, 0.1360, -0.1124, -1.2031, -5.877, -24.2595, -384.9590\}$. Evidently, the Hessian matrix is not positive definite, which causes the nonconvergence. In this case we will need to apply the relaxation method of equations (10)-(13) to get the solution

$$X = \begin{pmatrix} -3.0208 & -0.9591 & 2.1478 & 2.2488 & 0.9963 \\ -0.9591 & -0.8385 & 1.0214 & 0.6773 & 0.0265 \\ 2.1478 & 1.0214 & -2.4771 & -2.0446 & -1.1554 \\ 2.2488 & 0.6673 & -2.0446 & -2.3064 & -1.3623 \\ 0.9963 & 0.0265 & -1.1554 & -1.3623 & -1.5591 \end{pmatrix}$$

SUMMARY

In this note we have presented a fast numerical method for solving the algebraic Riccati matrix equation. In the example we investigated, the Newton method converged within a maximum of 5 iteration steps whatever the initial conditions. The merit of our method lies in the fact that it does not require additional assumption for the underlying Linear-Quadratic control model and therefore is capable of solving a wide range of applications. Furthermore, the derivative information used in the method will generally ensure fast convergence.

³Calculations were done in GAUSS on a 486 computer.

⁴This matrix was randomly selected

REFERENCES

- Amman, H.M., *Numerical Methods for Linear-Quadratic Models*, Handbook of Computational Economics (H.M. Amman, D.A. Kendrick and J. Rust, eds.), North-Holland, Amsterdam, Amsterdam, 1995.
- Amman, H.M. and D.A. Kendrick, *Nonconvexities in Stochastic Control Models*, International Economic Review **36** (1995), 455-475.
- Bertsekas, D.P., *Dynamic Programming and Stochastic Control*, Academic Press, New York, 1976.
- Kendrick, D.A., *Stochastic Control for Economic Models*, McGraw-Hill, New York, 1981.
- Kwakernaak, H. and R. Sivan, *Linear Optimal Control Systems*, John Wiley & Sons, New York, 1972.
- Lainiotis, D.G., N.D. Assimakis and S.K. Katsikas, *A New Computationally Effective Algorithm for Solving the Discrete Riccati Equation*, Journal of Mathematical Analysis **186** (1994), 868-895.
- Lancaster, P. and L. Rodman, *Algebraic Riccati Equations*, Clarendon Press, Oxford, 1995.
- Magnus, J.R. and H. Neudecker, *Matrix Differential Calculus with Applications in Statistics and Econometrics*, John Wiley, New York, 1994.
- McGratten, E.R., *A note on Computing Competitive Equilibria in Linear Models*, Journal of Economic Dynamics and Control **18** (1994), 149-160.
- Press, W.H., B.P. Flannery, S.A. Teukolsky and W.T. Vetterling, *Numerical Recipes 2nd edition*, Cambridge University Press, Cambridge, 1986.
- Ralston, A. and P. Rabinowitz, *A First Course in Numerical Analysis*, McGraw-Hill, Singapore, 1986.
- Sargent, T.J., *Dynamic Macroeconomic Theory*, Harvard University Press, Cambridge, 1987.
- Stoorvogel, A.A. and A.J.T.M. Weeren, *The Discrete-Time Riccati Equation Related to the H_∞ Control Problem*, IEEE Transactions on Automatic Control **39** (1994).
- Vaughan, D.R., *A Nonrecursive Algebraic Solution for the Riccati Equation*, IEEE Transactions on Automatic Control **15** (1970), 597-599.