

COVARIANT COMPLETELY POSITIVE LINEAR MAPS BETWEEN LOCALLY C^* -ALGEBRAS

MARIA JOIȚA

ABSTRACT. We prove a covariant version of the KSGNS (Kasparov, Stinespring, Gel'fand, Naimark, Segal) construction for completely positive linear maps between locally C^* -algebras. As an application of this construction, we show that a covariant completely positive linear map ρ from a locally C^* -algebra A to another locally C^* -algebra B with respect to a locally C^* -dynamical system (G, A, α) extends to a completely positive linear map on the crossed product $A \rtimes_\alpha G$.

1. INTRODUCTION

Locally C^* -algebras are generalizations of C^* -algebras. Instead of being given by a single norm, the topology on a locally C^* -algebra is defined by a directed family of C^* -seminorms. Such important concepts as Hilbert C^* -modules, adjointable operators, (completely) positive linear maps, C^* -dynamical systems can be defined with obvious modifications in the framework of locally C^* -algebras. The proofs are not always straightforward.

It is well-known that a positive linear functional on a C^* -algebra A induces a representation of this C^* -algebra on a Hilbert space by the GNS (Gel'fand, Naimark, Segal) construction (see, for example, [1]). Stinespring [13] extends this construction for completely positive linear map from A to $L(H)$, the C^* -algebra of all bounded linear operators on a Hilbert space H . On the other hand, Paschke [8] (respectively, Kasparov [5]) shows that a completely positive linear map from A to another C^* -algebra B (respectively, from A to the C^* -algebra of all adjointable operators on the Hilbert C^* -module H_B) induces a representation of A on a Hilbert B -module. In [2], the author extends the KSGNS (Kasparov, Stinespring, Gel'fand, Naimark, Segal) construction for a strict continuous, completely positive linear map

2000 Mathematical Subject Classification: 46L05, 46L08, 46L40

Keywords: locally C^* -algebras, Hilbert modules over locally C^* -algebras, locally C^* -dynamical system, covariant completely positive linear maps, KGNS (Kasparov, Stinespring, Gel'fand, Naimark, Segal) construction

This research was supported by grant CNCSIS (Romanian National Council for Research in High Education)-code A 1065/2006.

from a locally C^* -algebra A to $L_B(E)$, the locally C^* -algebra of all adjointable operators on a Hilbert module E over a locally C^* -algebra B . In this paper we propose to prove a covariant version of this construction. Thus we show that a covariant completely positive linear map from A to $L_B(E)$ with respect to a locally C^* -dynamical system (G, A, α) induces a non-degenerate, covariant representation of (G, A, α) on a Hilbert B -module which is unique up to unitary equivalence, Theorem 3.6. Using the analog of the covariant version of Stinespring construction [9] for bounded operators on Hilbert C^* -modules, Kaplan [4] shows that a discrete covariant completely positive map ρ from a unital C^* -algebra A to another unital C^* -algebra B extends to a completely positive map from the crossed product $A \rtimes_\alpha G$ to B . We extend this result showing that a non-degenerate, covariant, continuous completely positive linear map from a locally C^* -algebra to another locally C^* -algebra B extends to a non-degenerate, continuous completely positive linear map on the crossed product $A \rtimes_\alpha G$, Proposition 3.9.

2. PRELIMINARIES

A locally C^* -algebra is a complete complex Hausdorff topological $*$ -algebra whose topology is determined by a directed family of C^* -seminorms. If A is a locally C^* -algebra and $S(A)$ is the set of all continuous C^* -seminorms on A , then for each $p \in S(A)$, $A_p = A/\ker p$ is a C^* -algebra in the norm induced by p , and $\{A_p; \pi_{pq}\}_{p,q \in S(A), p \geq q}$, where π_{pq} is the canonical morphism from A_p onto A_q defined by $\pi_{pq}(a + \ker p) = a + \ker q$ for all $a \in A$ is an inverse system of C^* -algebras. Moreover, A can be identified with $\varprojlim_p A_p$. The canonical map from A onto A_p is denoted by π_p .

An approximate unit of A is an increasing net $\{e_\lambda\}_{\lambda \in \Lambda}$ of positive elements of A such that $p(e_\lambda) \leq 1$ for all $p \in S(A)$ and for all $\lambda \in \Lambda$, $p(ae_\lambda - a) \rightarrow 0$ and $p(e_\lambda a - a) \rightarrow 0$ for all $p \in S(A)$ and for all $a \in A$. Any locally C^* -algebra has an approximate unit [11, Proposition 3.11].

A morphism of locally C^* -algebras is a continuous $*$ -morphism from a locally C^* -algebra A to another locally C^* -algebra B . An isomorphism of locally C^* -algebras from A to B is a bijective map $\Phi : A \rightarrow B$ such that Φ and Φ^{-1} are morphisms of locally C^* -algebras.

Let $M_n(A)$ denote the $*$ -algebra of all $n \times n$ matrices over A with the algebraic operations and the topology obtained by regarding it as a direct sum of n^2 copies of A . Then $\{M_n(A_p); \pi_{pq}^{(n)}\}_{p,q \in S(A), p \geq q}$, where $\pi_{pq}^{(n)}([\pi_p(a_{ij})]_{i,j=1}^n) = [\pi_q(a_{ij})]_{i,j=1}^n$, is an inverse system of C^* -algebras and $M_n(A)$ can be identified with $\varprojlim_p M_n(A_p)$.

A linear map $\rho : A \rightarrow B$ between two locally C^* -algebras is completely positive if the linear maps $\rho^{(n)} : M_n(A) \rightarrow M_n(B)$ defined by

$$\rho^{(n)}([a_{ij}]_{i,j=1}^n) = [\rho(a_{ij})]_{i,j=1}^n$$

$n = 1, 2, \dots, n, \dots$, are all positive.

Definition 2.1. A *pre-Hilbert A -module* is a complex vector space E which is also a right A -module, compatible with the complex algebra structure, equipped with an A -valued inner product $\langle \cdot, \cdot \rangle : E \times E \rightarrow A$ which is $\mathbb{C}$ - and A -linear in its second variable and satisfies the following relations:

- (i) $\langle \xi, \eta \rangle^* = \langle \eta, \xi \rangle$ for every $\xi, \eta \in E$;
- (ii) $\langle \xi, \xi \rangle \geq 0$ for every $\xi \in E$;
- (iii) $\langle \xi, \xi \rangle = 0$ if and only if $\xi = 0$.

We say that E is a Hilbert A -module if E is complete with respect to the topology determined by the family of seminorms $\{\|\cdot\|_p\}_{p \in S(A)}$ where $\|\xi\|_p = \sqrt{p(\langle \xi, \xi \rangle)}$, $\xi \in E$ [11, Definition 4.1].

Let E be a Hilbert A -module. For $p \in S(A)$, $\mathcal{E}_p = \{\xi \in E; p(\langle \xi, \xi \rangle) = 0\}$ is a closed submodule of E and $E_p = E/\mathcal{E}_p$ is a Hilbert A_p -module with $(\xi + \mathcal{E}_p)\pi_p(a) = \xi a + \mathcal{E}_p$ and $\langle \xi + \mathcal{E}_p, \eta + \mathcal{E}_p \rangle = \pi_p(\langle \xi, \eta \rangle)$. The canonical map from E onto E_p is denoted by σ_p . For $p, q \in S(A)$, $p \geq q$ there is a canonical morphism of vector spaces σ_{pq} from E_p onto E_q such that $\sigma_{pq}(\sigma_p(\xi)) = \sigma_q(\xi)$, $\xi \in E$. Then $\{E_p; A_p; \sigma_{pq}\}_{p, q \in S(A), p \geq q}$ is an inverse system of Hilbert C^* -modules in the following sense: $\sigma_{pq}(\xi_p a_p) = \sigma_{pq}(\xi_p) \pi_{pq}(a_p)$, $\xi_p \in E_p, a_p \in A_p$; $\langle \sigma_{pq}(\xi_p), \sigma_{pq}(\eta_p) \rangle = \pi_{pq}(\langle \xi_p, \eta_p \rangle)$, $\xi_p, \eta_p \in E_p$; $\sigma_{pp}(\xi_p) = \xi_p$, $\xi_p \in E_p$ and $\sigma_{qr} \circ \sigma_{pq} = \sigma_{pr}$ if $p \geq q \geq r$, and $\lim_{\leftarrow p} E_p$ is a Hilbert A -module which can be identified with E [11, Proposition 4.4].

Let E and F be Hilbert A -modules. We say that an A -module morphism $T : E \rightarrow F$ is adjointable if there is an A -module morphism $T^* : F \rightarrow E$ such that $\langle T\xi, \eta \rangle = \langle \xi, T^*\eta \rangle$ for every $\xi \in E$ and $\eta \in F$. Any adjointable A -module morphism is continuous. The set $L_A(E, F)$ of all adjointable A -module morphisms from E into F becomes a locally convex space with topology defined by the family of seminorms $\{\tilde{p}\}_{p \in S(A)}$, where $\tilde{p}(T) = \|(\pi_p)_*(T)\|_{L_{A_p}(E_p, F_p)}$, $T \in L_A(E, F)$ and $(\pi_p)_*(T)(\xi + \mathcal{E}_p) = T\xi + \mathcal{F}_p$, $\xi \in E$. Moreover, $\{L_{A_p}(E_p, F_p); (\pi_{pq})_*\}_{p, q \in S(A), p \geq q}$, where $(\pi_{pq})_* : L_{A_p}(E_p, F_p) \rightarrow L_{A_q}(E_q, F_q)$, $(\pi_{pq})_*(T_p)(\sigma_q(\xi)) = \chi_{pq}(T_p(\sigma_p(\xi)))$, and χ_{pq} , $p, q \in S(A)$, $p \geq q$ are the connecting maps of the inverse system $\{F_p\}_{p \in S(A)}$, is an inverse system of Banach spaces, and $\lim_{\leftarrow p} L_{A_p}(E_p, F_p)$ can be identified with $L_A(E, F)$ [11, Proposition 4.7]. Thus topologized, $L_A(E, E)$ becomes a locally C^* -algebra, and we write $L_A(E)$ for $L_A(E, E)$.

The strict topology on $L_A(E)$ is defined by the family of seminorms $\{\|\cdot\|_{p,\xi}\}_{(p,\xi) \in S(A) \times E}$, where $\|T\|_{p,\xi} = \|T\xi\|_p + \|T^*\xi\|_p$, $T \in L_A(E)$.

Two Hilbert A -modules E and F are unitarily equivalent if there is a unitary element in $L_A(E, F)$.

A non-degenerate representation of a locally C^* -algebra A on a Hilbert module E over a locally C^* -algebra B is a morphism of locally C^* -algebras Φ from A to $L_B(E)$ such that $\Phi(A)E$ is dense in E .

A continuous completely positive linear map ρ from A to $L_B(E)$ is non-degenerate if the net $\{\rho(e_\lambda)\}_{\lambda \in \Lambda}$ converges strictly to the identity map on E for some approximate unit $\{e_\lambda\}_{\lambda \in \Lambda}$ for A .

Let G be a locally compact group and let A be a locally C^* -algebra. An action of G on A is a morphism α from G to $\text{Aut}(A)$, the set of all isomorphisms of locally C^* -algebras from A to A . The action α is continuous if the function $(t, a) \rightarrow \alpha_t(a)$ from $G \times A$ to A is jointly continuous. An action α is called an inverse limit action if we can write A as inverse limit $\varprojlim_{\delta} A_\delta$ of C^* -algebras in such a way that there are actions $\alpha^{(\delta)}$ of G on A_δ such that $\alpha_t = \varprojlim_{\delta} \alpha_t^{(\delta)}$ for all t in G [12, Definition 5.1]. An action α of G on A is a continuous inverse limit action if there is a cofinal subset $S_G(A, \alpha)$ of G -invariant continuous C^* -seminorms on A (a continuous C^* -seminorm p on A is G -invariant if $p(\alpha_t(a)) = p(a)$ for all a in A and for all t in G). So if α is a continuous inverse limit action of G on A we can suppose that $S(A) = S_G(A, \alpha)$.

A locally C^* -dynamical system is a triple (G, A, α) , where G is a locally compact group, A is a locally C^* -algebra and α is a continuous action of G on A .

Let α be a continuous inverse limit action of G on A . The set $C_c(G, A)$ of all continuous functions from G to A with compact support becomes a $*$ -algebra with convolution of two functions

$$(f \times h)(s) = \int_G f(t) \alpha_t(h(t^{-1}s)) dt$$

as product and involution defined by

$$f^\sharp(t) = \Delta(t)^{-1} \alpha_t(f(t^{-1})^*)$$

where Δ is the modular function on G . The Hausdorff completion of $C_c(G, A)$ with respect to the topology defined by the family of submultiplicative $*$ -seminorms $\{N_p\}_{p \in S(A)}$, where

$$N_p(f) = \int_G p(f(s)) ds$$

is denoted by $L^1(G, A, \alpha)$ and the enveloping locally C^* -algebra $A \rtimes_\alpha G$ of $L^1(G, A, \alpha)$ is called the crossed product of A by α [3, Definition 3.14]. Moreover, the C^* -algebras $(A \rtimes_\alpha G)_p$ and $A_p \rtimes_{\alpha(p)} G$ can be identified for each $p \in S(A)$ and so $A \rtimes_\alpha G$ can be identified with $\lim_{\leftarrow p} A_p \rtimes_{\alpha(p)} G$ [3, Remark 3.15].

3. COVARIANT REPRESENTATIONS ASSOCIATED WITH A COVARIANT COMPLETELY POSITIVE LINEAR MAP

Let B be a locally C^* -algebra, let E be a Hilbert B -module and let G be a locally compact group.

Definition 3.1. A unitary representation of G on E is a map u from G to $L_B(E)$ such that

- (a) u_g is a unitary element in $L_B(E)$ for all $g \in G$;
- (b) $u_{gt} = u_g u_t$ for all $g, t \in G$;
- (c) the map $g \mapsto u_g \xi$ from G to E is continuous for all $\xi \in E$.

Remark 3.2. If u is a unitary representation of G on E , then for each $q \in S(B)$, $g \mapsto (\pi_q)_* \circ u$ is a unitary representation of G on E_q . Moreover, $u_g = \lim_{\leftarrow q} u_g^{(q)}$, where $u_g^{(q)} = (\pi_q)_*(u_g)$, for all $g \in G$.

Definition 3.3. A non-degenerate, covariant representation of a locally C^* -dynamical system (G, A, α) on a Hilbert B -module E is a triple (Φ, v, E) , where Φ is a non-degenerate representation of A on E , v is a unitary representation of G on E and

$$\Phi(\alpha_g(a)) = v_g \Phi(a) v_g^*$$

for all $g \in G$ and $a \in A$.

Proposition 3.4. Let (G, A, α) be a locally C^* -dynamical system such that α is an inverse limit action, let (Φ, v, E) be a non-degenerate covariant representation of (G, A, α) on a Hilbert B -module E . Then there is a unique non-degenerate representation $\Phi \rtimes v$ of the crossed product $A \rtimes_\alpha G$ on E such that

$$(\Phi \rtimes v)(f) = \int_G \Phi(f(g)) v_g dg$$

for all $f \in C_c(G, A)$.

PROOF. We partition the proof into two steps.

Step1. We suppose that B is a C^* -algebra.

Since Φ is a representation of A on E , there is $p \in S(A)$ such that $\|\Phi(a)\|_{L_B(E)} \leq p(a)$ for all $a \in A$. From this fact, we deduce that there is a morphism of C^* -algebras Φ_p from A_p to $L_B(E)$ such that $\Phi_p \circ \pi_p = \Phi$. Therefore Φ_p is a representation of A_p on E , and moreover, it is non-degenerate, since Φ is non-degenerate and π_p is surjective. It is not difficult to check that (Φ_p, v, E) is a non-degenerate covariant representation of $(G, A_p, \alpha^{(p)})$. Then there is a unique non-degenerate representation $\Phi_p \times v$ of $A_p \times_{\alpha^{(p)}} G$ on E such that

$$(\Phi_p \times v)(f) = \int_G \Phi_p(f(g))v_g dg$$

for all $f \in C_c(G, A_p)$ (see, for example, Proposition 7.6.4, [10]). Therefore $\Phi \times v = (\Phi_p \times v) \circ \tilde{\pi}_p$, where $\tilde{\pi}_p$ is the canonical map from $A \times_\alpha G$ onto $A_p \times_{\alpha^{(p)}} G$ is a non-degenerate representation of $A \times_\alpha G$ on E such that

$$(\Phi \times v)(f) = \int_G \Phi_p(\tilde{\pi}_p(f)(g))v_g dg = \int_G \Phi(f(g))v_g dg$$

for all $f \in C_c(G, A)$, and since $C_c(G, A)$ is dense in $A \times_\alpha G$, $\Phi \times v$ is unique with the above property.

Step2. The general case.

For each $q \in S(B)$, $(\pi_q)_* \circ \Phi$ is a non-degenerate representation of A on E_q , and $((\pi_q)_* \circ \Phi, v^{(q)}, E_q)$ is a non-degenerate covariant representation of (G, A, α) on E_q . By Step 1 there is a unique non-degenerate representation $((\pi_q)_* \circ \Phi) \times v^{(q)}$ of $A \times_\alpha G$ on E_q such that

$$(((\pi_q)_* \circ \Phi) \times v^{(q)})(f) = \int_G (\pi_q)_*(\Phi(f(g)))v_g^{(q)} dg$$

for all $f \in C_c(G, A)$. By Lemma 3.7 in [3], we have

$$\begin{aligned} (\pi_{qr})_*(((\pi_q)_* \circ \Phi) \times v^{(q)})(f) &= \int_G (\pi_{qr})_*((\pi_q)_*(\Phi(f(g))))v_g^{(q)} dg \\ &= \int_G (\pi_r)_*(\Phi(f(g)))v_g^{(r)} dg \\ &= (((\pi_r)_* \circ \Phi) \times v^{(r)})(f) \end{aligned}$$

for all $f \in C_c(G, A)$ and for all $q, r \in S(B)$ with $q \geq r$. Therefore $(\pi_{qr})_* \circ (((\pi_q)_* \circ \Phi) \times v^{(q)}) = ((\pi_r)_* \circ \Phi) \times v^{(r)}$ for all $q, r \in S(B)$ with $q \geq r$. This implies that there is a continuous $*$ -morphism $\Phi \times v$ from $A \times_\alpha G$ to $L_B(E)$ such that $(\pi_q)_* \circ (\Phi \times v) = ((\pi_q)_* \circ \Phi) \times v^{(q)}$ for all $q \in S(B)$. Using Lemma III 3.1 in [7],

it is not hard to check that $(\Phi \times v)(A \times_\alpha G)E$ is dense in E . Therefore $\Phi \times v$ is a non-degenerate representation of $A \times_\alpha G$ on E , and by Lemma 3.7 in [3],

$$(\Phi \times v)(f) = \int_G \Phi(f(g))v_g dg$$

for all $f \in C_c(G, A)$. Moreover, since $C_c(G, A)$ is dense in $A \times_\alpha G$, $\Phi \times v$ is unique with the above property. q.e.d.

Definition 3.5. Let (G, A, α) be a locally C^* -dynamical system and let u be a unitary representation of G on a Hilbert B -module E . We say that a completely positive linear map ρ from A to $L_B(E)$ is u -covariant with respect to the locally C^* -dynamical system (G, A, α) if

$$\rho(\alpha_g(a)) = u_g \rho(a) u_g^*$$

for all $a \in A$ and for all $g \in G$.

Recall that if ρ is a completely positive linear map from a C^* -algebra A to $L_B(E)$, the C^* -algebra of all adjointable operators on a Hilbert module E over a C^* -algebra B , the quotient vector space $(A \otimes_{\text{alg}} E) / \mathcal{N}_\rho$, where $\mathcal{N}_\rho = \{\sum_{i=1}^n a_i \otimes \xi_i; \sum_{i,j=1}^n \langle \xi_i, \rho(a_i^* a_j) \xi_j \rangle = 0\}$ becomes a pre-Hilbert B -module with the action of B on $A \otimes_{\text{alg}} E$ defined by $(a \otimes \xi + \mathcal{N}_\rho) b = a \otimes \xi b + \mathcal{N}_\rho$ and the inner-product defined by

$$\left\langle \sum_{i=1}^n a_i \otimes \xi_i + \mathcal{N}_\rho, \sum_{j=1}^m b_j \otimes \eta_j + \mathcal{N}_\rho \right\rangle_\rho = \sum_{i=1}^n \sum_{j=1}^m \langle \xi_i, \rho(a_i^* b_j) \eta_j \rangle.$$

The following theorem is a covariant version of Theorem 4.6 in [2].

Theorem 3.6. Let (G, A, α) be a locally C^* -dynamical system, let u be a unitary representation of G on a Hilbert module E over a locally C^* -algebra B , and let ρ be a u -covariant, non-degenerate, continuous completely positive linear map from A to $L_B(E)$.

1. Then there is a covariant representation $(\Phi_\rho, v^\rho, E_\rho)$ of (G, A, α) and an element V_ρ in $L_B(E, E_\rho)$ such that
 - (a) $\rho(a) = V_\rho^* \Phi_\rho(a) V_\rho$ for all $a \in A$;
 - (b) $\{\Phi_\rho(a) V_\rho \xi; a \in A, \xi \in E\}$ spans a dense submodule of E_ρ ;
 - (c) $v_g^\rho V_\rho = V_\rho u_g$ for all $g \in G$.
2. If F is a Hilbert B -module, (Φ, v, F) is a covariant representation of (G, A, α) and W is an element in $L_B(E, F)$ such that
 - (a) $\rho(a) = W^* \Phi(a) W$ for all $a \in A$;
 - (b) $\{\Phi(a) W \xi; a \in A, \xi \in F\}$ spans a dense submodule of F ;

(c) $v_g W = W u_g$ for all $g \in G$,

then there is a unitary operator U in $L_B(E_\rho, F)$ such that

- (i) $\Phi(a)U = U\Phi_\rho(a)$ for all $a \in A$;
- (ii) $v_g U = U v_g^\rho$ for all $g \in G$;
- (iii) $W = UV_\rho$.

PROOF. We partition the proof into two steps.

Step 1. Suppose that B is a C^* -algebra.

1. Let $\{e_\lambda\}_{\lambda \in \Lambda}$ be an approximate unit of A such that the net $\{\rho(e_\lambda)\}_{\lambda \in \Lambda}$ is strictly convergent to the identity operator on E , and let $(\Phi_\rho; V_\rho; E_\rho)$ be the KSGNS construction associated with ρ . Since ρ is continuous there is $p \in S(A)$ and a completely positive linear map ρ_p from A_p to $L_B(E)$ such that $\rho = \rho_p \circ \pi_p$ (see, for example, the proof of Proposition 3.5 in [2]). By the proof of Theorem 4.6 in [2] we can suppose that E_ρ is the completion of the pre-Hilbert space $(A_p \otimes_{\text{alg}} E) / \mathcal{N}_{\rho_p}$, $V_\rho \xi = \lim_{\lambda} (\pi_p(e_\lambda) \otimes \xi + \mathcal{N}_{\rho_p})$ and $\Phi_\rho(a) (\pi_p(b) \otimes \xi + \mathcal{N}_{\rho_p}) = \pi_p(ab) \otimes \xi + \mathcal{N}_{\rho_p}$ for all $a, b \in A$ and for all $\xi \in E$.

Let $g \in G$. From

$$\begin{aligned} & \left\langle \sum_{i=1}^n \pi_p(a_i) \otimes \xi_i + \mathcal{N}_{\rho_p}, \sum_{j=1}^m b_j \otimes \eta_j + \mathcal{N}_{\rho_p} \right\rangle_{\rho_p} = \sum_{i=1}^n \sum_{j=1}^m \langle \xi_i, \rho(a_i^* b_j) \eta_j \rangle \\ &= \sum_{i=1}^n \sum_{j=1}^m \langle u_g(\xi_i), u_g \rho(a_i^* b_j) u_{g^{-1}} u_g(\eta_j) \rangle \\ &= \sum_{i=1}^n \sum_{j=1}^m \langle u_g(\xi_i), \rho(\alpha_g(a_i^* b_j)) u_g(\eta_j) \rangle \\ &= \left\langle \sum_{i=1}^n \pi_p(\alpha_g(a_i)) \otimes u_g(\xi_i) + \mathcal{N}_{\rho_p}, \sum_{j=1}^m \pi_p(\alpha_g(b_j)) \otimes u_g(\eta_j) + \mathcal{N}_{\rho_p} \right\rangle_{\rho_p} \end{aligned}$$

for all $\xi_1, \dots, \xi_n, \eta_1, \dots, \eta_m \in E$, for all $a_1, \dots, a_n, b_1, \dots, b_m \in A$ and for all $g \in G$, we deduce that, there is a unitary operator v_g^ρ in $L_B(E_\rho)$ such that

$$v_g^\rho(\pi_p(a) \otimes \xi + \mathcal{N}_{\rho_p}) = \pi_p(\alpha_g(a)) \otimes u_g \xi + \mathcal{N}_{\rho_p}$$

for all $a \in A$ and for all $\xi \in E$. It is not difficult to check that the map $g \mapsto v_g^\rho$ from G to $L_B(E_\rho)$ is a unitary representation of G on E_ρ .

To show that $(\Phi_\rho, v^\rho, E_\rho)$ is a covariant representation of (G, A, α) it remains to prove that $\Phi_\rho(\alpha_g(a)) = v_g^\rho \Phi_\rho(a) v_{g^{-1}}^\rho$ for all $g \in G$ and $a \in A$. Let $g \in G$ and $a \in A$. We have

$$\begin{aligned}
(v_g^\rho \Phi_\rho(a) v_{g^{-1}}^\rho)(\pi_p(b) \otimes \xi + \mathcal{N}_{\rho_p}) &= (v_g^\rho \Phi_\rho(a))(\pi_p(\alpha_{g^{-1}}(b)) \otimes u_{g^{-1}}\xi + \mathcal{N}_{\rho_p}) \\
&= v_g^\rho(\pi_p(a\alpha_{g^{-1}}(b)) \otimes u_{g^{-1}}\xi + \mathcal{N}_{\rho_p}) \\
&= \pi_p(\alpha_g(a)b) \otimes \xi + \mathcal{N}_{\rho_p} \\
&= (\Phi_\rho(\alpha_g(a)))(\pi_p(b) \otimes \xi + \mathcal{N}_{\rho_p})
\end{aligned}$$

for all $b \in A$ and for all $\xi \in E$. Hence $\Phi_\rho(\alpha_g(a)) = v_g^\rho \Phi_\rho(a) v_{g^{-1}}^\rho$.

By Theorem 4.6 (1), [2] the conditions (a) and (b) are verified. To show that the condition (c) is verified, let $\xi \in E$ and $g \in G$. Then we have

$$\begin{aligned}
\|v_g^\rho V_\rho \xi - V_\rho u_g \xi\|^2 &= \lim_{\lambda \in \Lambda} \left\| v_g^\rho \left(\pi_p(e_\lambda) \otimes \xi + \mathcal{N}_{\rho_p} \right) - V_\rho u_g \xi \right\|^2 \\
&= \lim_{\lambda \in \Lambda} \left\| \langle \xi, \rho(e_\lambda^2) \xi \rangle + \langle \xi, \xi \rangle - \langle \rho(\alpha_g(e_\lambda)) u_g \xi, u_g \xi \rangle - \langle u_g \xi, \rho(\alpha_g(e_\lambda)) u_g \xi \rangle \right\| \\
&\leq \lim_{\lambda \in \Lambda} \left\| \langle \xi, \rho(e_\lambda) \xi \rangle + \langle \xi, \xi \rangle - \langle \rho(e_\lambda) \xi, \xi \rangle - \langle \xi, \rho(e_\lambda) \xi \rangle \right\| \\
&= \lim_{\lambda \in \Lambda} \left\| \langle \xi - \rho(e_\lambda) \xi, \xi \rangle \right\| = 0.
\end{aligned}$$

Therefore the condition (c) is also verified.

2. By Theorem 4.6 (2), [2] there is a unitary operator U in $L_B(E_\rho, F)$ defined by $U(\Phi_\rho(a) V_\rho \xi) = \Phi(a) W \xi$ such that $\Phi(a) U = U \Phi_\rho(a)$ for all $a \in A$, and $W = U V_\rho$.

Let $g \in G$. From

$$\begin{aligned}
(v_g U)(\Phi_\rho(a) V_\rho \xi) &= v_g(\Phi(a) W \xi) = \Phi(\alpha_g(a)) v_g W \xi \\
&= \Phi(\alpha_g(a)) W u_g \xi = U(\Phi_\rho(\alpha_g(a)) V_\rho u_g \xi) \\
&= U(\Phi_\rho(\alpha_g(a)) v_g^\rho V_\rho \xi) = (U v_g^\rho)(\Phi_\rho(a) V_\rho \xi).
\end{aligned}$$

for all $a \in A$ and for all $\xi \in E$, we conclude that $v_g U = U v_g^\rho$ and thus the assertion 2. is proved.

Step 2. The general case.

Let $q \in S(B)$. Then $\rho_q = (\pi_q)_* \circ \rho$ is a $u^{(q)}$ -covariant, non-degenerate, continuous completely positive linear map from A to $L_{B_q}(E_q)$, $((\pi_q)_* \circ \Phi, v^{(q)}, F_q)$ is a covariant representation of (G, A, α) and $(\pi_q)_*(W)$ is an element in $L_{B_q}(E_q, F_q)$ such that the conditions (a), (b) and (c) from 2. are verified. By Step 1, there is a covariant representation $(\Phi_{\rho_q}, v^{\rho_q}, E_{\rho_q})$ of (G, A, α) and an element V_{ρ_q} in $L_{B_q}(E_q, E_{\rho_q})$ which verify the conditions (a), (b) and (c) from 1. and there is a unitary operator U_q in $L_{B_q}(E_{\rho_q}, F_q)$ which verifies the conditions i), ii) and iii) from 2.

Let $(\Phi_\rho; V_\rho; E_\rho)$ be the KSGNS construction associated with ρ . According to the proof of Theorem 4.6 in [2], $(\pi_q)_* \circ \Phi_\rho = \Phi_{\rho_q}$; $(\pi_q)_*(V_\rho) = V_{\rho_q}$; $(E_\rho)_q = E_{\rho_q}$ for all $q \in S(B)$ and $(U_q)_q$ is a coherent sequence in $L_{B_q}(E_{\rho_q}, F_q)$. It is not difficult to check that for each $g \in G$, $(v_g^{\rho_q})_q$ is a coherent sequence in $L_{B_q}(E_{\rho_q})$, and the map $g \mapsto v_g^\rho$, where v_g^ρ is an element in $L_B(E_\rho)$ such that $(\pi_q)_*(v_g^\rho) = v_g^{\rho_q}$ for all

$q \in S(B)$ is a unitary representation of G on E_ρ . Also it is not difficult to check that $(\Phi_\rho, v^\rho, E_\rho)$ is a covariant representation of (G, A, α) which verifies the conditions (a), (b) and (c) from 1.

Let $U \in L_B(E_\rho, F)$ such that $(\pi_q)_*(U) = U_q$ for all $q \in S(B)$. Clearly U is a unitary operator in $L_B(E_\rho, F)$ and it verifies the conditions *i*), *ii*) and *iii*) from 2. q.e.d.

Remark 3.7. *The covariant representation $(\Phi_\rho, v^\rho, E_\rho)$ of (G, A, α) induced by ρ is unique up to unitary equivalence.*

From Proposition 3.4 and Theorem 3.6 we obtain the following corollary.

Corollary 3.8. *Let (G, A, α) be a locally C^* -dynamical system such that α is an inverse limit action, let u be a unitary representation of G on a Hilbert module E over a locally C^* -algebra B , and let ρ be a u -covariant, non-degenerate, continuous completely positive linear map from A to $L_B(E)$. Then ρ induces a non-degenerate representation of the crossed product $A \times_\alpha G$ on a Hilbert B -module.*

The following proposition is a generalization of Proposition 2 in [4] in the context of locally C^* -algebras.

Proposition 3.9. *Let (G, A, α) be a locally C^* -dynamical system such that α is an inverse limit action, let B be a locally C^* -algebra, let E be a Hilbert B -module and let u be a unitary representation of G on E . If ρ is a u -covariant, non-degenerate, continuous completely positive linear map from A to $L_B(E)$, then there is a unique completely positive linear map φ from $A \times_\alpha G$ to $L_B(E)$ such that*

$$\varphi(f) = \int_G \rho(f(g)) u_g dg$$

for all $f \in C_c(G, A)$. Moreover, φ is non-degenerate.

PROOF. By Theorem 3.6 there is a covariant representation $(\Phi_\rho, v^\rho, E_\rho)$ of (G, A, α) and an element V_ρ in $L_B(E, E_\rho)$ such that $\rho(a) = V_\rho^* \Phi_\rho(a) V_\rho$ and $v_g^\rho V_\rho = V_\rho u_g$ for all $a \in A$, and for all $g \in G$.

Let $\Phi_\rho \times v^\rho$ be the representation of $A \times_\alpha G$ associated with $(\Phi_\rho, v^\rho, E_\rho)$. We define $\varphi : A \times_\alpha G$ to $L_B(E)$ by

$$\varphi(x) = V_\rho^* (\Phi_\rho \times v^\rho)(x) V_\rho.$$

Clearly φ is a continuous completely positive linear map from $A \times_\alpha G$ to $L_B(E)$. Let $\{e_\lambda\}_{\lambda \in \Lambda}$ be an approximate unit for $A \times_\alpha G$ and let $\xi \in E$. Since $\Phi_\rho \times v^\rho$ is

non-degenerate, by Proposition 4.2 in [2]

$$\lim_{\lambda} V_{\rho}^*(\Phi_{\rho} \times v^{\rho})(e_{\lambda})V_{\rho}\xi = V_{\rho}^*V_{\rho}\xi = \xi.$$

This implies that the net $\{\rho(e_{\lambda})\}_{\lambda \in \Lambda}$ converges strictly to the identity map on E , and so φ is non-degenerate.

For $f \in C_c(G, A)$ we have

$$\begin{aligned} \varphi(f) &= V_{\rho}^*(\Phi_{\rho} \times v^{\rho})(f)V_{\rho} = \int_G V_{\rho}^*\Phi_{\rho}(f(g))v_g^{\rho}V_{\rho}dg \\ &= \int_G V_{\rho}^*\Phi_{\rho}(f(g))V_{\rho}u_gdg = \int_G \rho(f(g))u_gdg \end{aligned}$$

and since $C_c(G, A)$ is dense in $A \times_{\alpha} G$, φ is unique with this property. q.e.d.

Corollary 3.10. *Let (G, A, α) be a locally C^* -dynamical system, let B be a locally C^* -algebra, let E be a Hilbert B -module, let u be a unitary representation of G on E , and let ρ be a u -covariant, non-degenerate, continuous completely positive linear map from A to $L_B(E)$. If G is a compact group, then there is a unique completely positive linear map φ from $A \times_{\alpha} G$ to $L_B(E)$ such that*

$$\varphi(f) = \int_G \rho(f(g))u_gdg$$

for all $f \in C_c(G, A)$. Moreover, φ is non-degenerate.

PROOF. The corollary follows from Proposition 3.9, since G is compact and then α is an inverse limit action of G on A [12, Lemma 5.2].q.e.d.

REFERENCES

- [1] M. Fragoulopoulou, *An introduction to the representation theory of topological $*$ -algebras*, Schriftenreihe, Univ. Münster, **48**(1988), 1-81.
- [2] M. Joița, *Strict completely positive maps between locally C^* -algebras and representations on Hilbert modules*, J. London Math. Soc. (2), **66**(2002), 421-432.
- [3] M. Joița, *Crossed products of locally C^* -algebras*, Rocky Mountain J. Math. (to appear).
- [4] A. Kaplan, *Covariant completely positive maps and liftings*, Rocky Mountain J. Math. **23**(1993), 939-946.
- [5] G. G. Kasparov, *Hilbert C^* -modules: Theorem of Stinespring and Voiculescu*, J. Operator Theory **4**(1980), 133-150.
- [6] E. C. Lance, *Hilbert C^* -modules. A toolkit for operator algebraists*, London Mathematical Society Lecture Note Series 210, Cambridge University Press, Cambridge 1995.
- [7] A. Mallios, *Topological algebras: Selected Topics*, North Holland, Amsterdam, 1986.
- [8] W. L. Paschke, *Inner product modules over B^* -algebras*, Trans. Amer. Math. Soc. **182**(1973), 443-468.
- [9] V. Paulsen, *A covariant version of Ext*, Michigan, Math. J. **29**(1982), 131-142.

- [10] G. K. Pedersen, *C*-algebras and their automorphism groups*, Academic Press, London, New-York, San Francisco, 1979.
- [11] N. C. Phillips, *Inverse limit of C*-algebras*, J. Operator Theory, **19**(1988), 159-195.
- [12] N. C. Phillips, *Representable K-theory for σ -C*-algebras*, K-Theory, **3**(1989),5, 441-478.
- [13] W. Stinespring, *Positive functions on C*-algebras*, Proc. Amer. Math. Soc., **6**(1955), 211-216.

Department of Mathematics, Faculty of Chemistry, University of Bucharest, Bd.
Regina Elisabeta nr.4-12, Bucharest, Romania
mjoita@fmi.unibuc.ro