

# Determining Distributions

Daniel A. Menascé, Ph.D.  
Dept of Computer Science  
George Mason University

© 2001 D. A. Menascé. All Rights Reserved.

1

## Problem

- Given a data set, determine the best distribution that fits the data.
- Useful for simulation and analytic studies.
- Start by plotting a histogram to narrow down type of distribution.
- Then, use a quantile-quantile plot.

© 2001 D. A. Menascé. All Rights Reserved.

2

## Quantile-Quantile plot

$y_i$ : observed  $q_i$ -th quantile

$x_i$ : theoretical  $q_i$ -th quantile

If the observations come from the same distribution, the quantile-quantile plot should be linear.

To determine the theoretical quantile, we need to be able to invert the CDF  $F(x)$ .

$$q_i = F(x_i)$$

$$x_i = F^{-1}(q_i)$$

© 2001 D. A. Menascé. All Rights Reserved.

3

## Examples of CDFs and Their Inverse Functions

Exponential	$F(x) = 1 - e^{-x/a}$	$-a \ln(1-u)$
Pareto	$F(x) = 1 - x^{-a}$	$\frac{1}{(1-u)^{1/a}}$
Geometric	$F(x) = 1 - (1-p)^x$	$\left\lceil \frac{\ln(u)}{\ln(1-p)} \right\rceil$

© 2001 D. A. Menascé. All Rights Reserved.


4

## Example of a Quantile-Quantile Plot

- One thousand values are suspected of coming from an exponential distribution (see histogram in the next slide). The quantile-quantile plot is pretty much linear, which confirms the conjecture.

© 2001 D. A. Menascé. All Rights Reserved.

5



© 2001 D. A. Menascé. All Rights Reserved.


6

## Data for Quantile-Quantile Plot

qi	yi	xi
0.100	0.22	0.21
0.200	0.49	0.45
0.300	0.74	0.71
0.400	1.03	1.02
0.500	1.41	1.39
0.600	1.84	1.83
0.700	2.49	2.41
0.800	3.26	3.22
0.900	4.31	4.61
0.930	4.98	5.32
0.950	5.49	5.99
0.970	6.53	7.01
0.980	7.84	7.82
0.985	8.12	8.40
0.990	8.82	9.21
1.000	17.91	18.42

© 2001 D. A. Menascé. All Rights Reserved.

7



© 2001 D. A. Menascé. All Rights Reserved.

8

## What if the Inverse of the CDF Cannot be Found?

- Use tables and interpolate.
- Approximation for  $N(0,1)$ :


$$x_i = 4.91[q_i^{0.14} - (1 - q_i)^{0.14}]$$

- For  $N(\mu, \sigma)$  the  $x_i$  values are scaled as

$$\mathbf{m} + \mathbf{S}x_i \quad \text{before plotting.}$$

© 2001 D. A. Menascé. All Rights Reserved.

9



© 2001 D. A. Menascé. All Rights Reserved.


10

## Using PhStat

- The normality assumption can be checked with PhStat by using the Normal Probability Plot option from the Probability Distribution option. (example in next slide).

© 2001 D. A. Menascé. All Rights Reserved.

11



© 2001 D. A. Menascé. All Rights Reserved.

12

