

Individual Events

SI	$A$	15	I1	$R$	30	I2	$a$	16	I3	$m$	3	I4	$m$	3
	$B$	3		$S$	120		$b$	$\frac{3}{2}$		$n$	9		$n$	$\frac{9}{4}$
	$C$	4		$T$	11		$c$	36		$p$	2		$p$	9
	$D$	8		$U$	72		$d$	42		$q$	1141		$q$	8

Group Events

SG	$z$	540	G1	$q$	3	G2	$A$	$-\frac{17}{13}$	G3	$A$	5	G4	$P$	$\frac{3}{8}$
	$R$	6		$k$	1		$B$	13		$R$	4018		$R$	$\frac{1}{2}$
	$k$	5		$w$	25		$C$	46		$Q$	$\frac{4\sqrt{5}}{5}$		$S$	320
	$xyz$	1		$p$	$\frac{3}{2}$		$D$	30		$T$	$5-2\sqrt{3}$		$Q$	-1

Sample Individual Event

SI.1 Let  $A = 15 \times \tan 44^\circ \times \tan 45^\circ \times \tan 46^\circ$ , find the value of  $A$ .

$$A = 15 \times \tan 44^\circ \times 1 \times \frac{1}{\tan 44^\circ} = 15$$

SI.2 Let  $n$  be a positive integer and  $\overbrace{20082008 \dots 2008}^{n \text{ 2008's}}$  is divisible by  $A$ . If the least possible value of  $n$  is  $B$ , find the value of  $B$ .

The given number is divisible by 15. Therefore it is divisible by 3 and 5.

The last 2 digits of the given number is 15, which is divisible by 15.

The necessary condition is:  $\overbrace{20082008 \dots 2008}^{n \text{ 2008's}}$  must be divisible by 3.

$2 + 0 + 0 + 8 = 10$  which is not divisible by 3.

The least possible  $n$  is 3:  $2+0+0+8+2+0+0+8+2+0+0+8 = 30$  which is divisible by 3.

SI.3 Given that there are  $C$  integers that satisfy the equation  $|x - 2| + |x + 1| = B$ , find the value of  $C$

$$|x - 2| + |x + 1| = 3$$

If  $x < -1$ ,  $2 - x - x - 1 = 3 \Rightarrow x = -1$  (rejected)

If  $-1 \leq x \leq 2$ ,  $2 - x + x + 1 = 3 \Rightarrow 3 = 3$ , always true  $-1 \leq x \leq 2$

If  $2 < x$ ,  $x - 2 + x + 1 = 3 \Rightarrow x = 2$  (reject)

$$-1 \leq x \leq 2 \text{ only}$$

$\therefore x$  is an integer,  $x = -1, 0, 1, 2$ ;  $C = 4$

SI.4 In the coordinate plane, the distance from the point  $(-C, 0)$  to the straight line  $y = x$  is  $\sqrt{D}$ , find the value of  $D$ .

The distance from  $P(x_0, y_0)$  to the straight line  $Ax + By + C = 0$  is  $\left| \frac{Ax_0 + By_0 + C}{\sqrt{A^2 + B^2}} \right|$ .

The distance from  $(-4, 0)$  to  $x - y = 0$  is  $\sqrt{D} = \left| \frac{-4 - 0 + 0}{\sqrt{1^2 + (-1)^2}} \right| = \frac{4}{\sqrt{2}} = 2\sqrt{2} = \sqrt{8}$ ;  $D = 8$

## Individual Event 1

11.1 Let  $a, b, c$  and  $d$  be the distinct roots of the equation  $x^4 - 15x^2 + 56 = 0$ .

If  $R = a^2 + b^2 + c^2 + d^2$ , find the value of  $R$ .

$$x^4 - 15x^2 + 56 = 0 \Rightarrow (x^2 - 7)(x^2 - 8) = 0$$

$$a = \sqrt{7}, b = -\sqrt{7}, c = \sqrt{8}, d = -\sqrt{8}$$

$$R = a^2 + b^2 + c^2 + d^2 = 7 + 7 + 8 + 8 = 30$$

11.2 In Figure 1,  $AD$  and  $BE$  are straight lines with  $AB = AC$  and  $AB \parallel ED$ .


If  $\angle ABC = R^\circ$  and  $\angle ADE = S^\circ$ , find the value of  $S$ .

$$\angle ABC = 30^\circ = \angle ACB \quad (\text{base } \angle \text{ isos. } \Delta)$$

$$\angle BAC = 120^\circ \quad (\angle \text{s sum of } \Delta)$$

$$\angle ADE = 120^\circ \quad (\text{alt. } \angle \text{s } AB \parallel ED)$$

$$S = 120$$



11.3 Let  $F = 1 + 2 + 2^2 + 2^3 + \dots + 2^S$  and  $T = \sqrt{\frac{\log(1+F)}{\log 2}}$ , find the value of  $T$ .

$$F = 1 + 2 + 2^2 + 2^3 + \dots + 2^{120} = \frac{2^{121} - 1}{2 - 1} = 2^{121} - 1$$

$$T = \sqrt{\frac{\log(1+F)}{\log 2}} = \sqrt{\frac{\log 2^{121}}{\log 2}} = 11$$

11.4 Let  $f(x)$  be a function such that  $f(n) = (n-1)f(n-1)$  and  $f(n) \neq 0$  hold for all integers  $n \geq 6$ .

If  $U = \frac{f(T)}{(T-1)f(T-3)}$ , find the value of  $U$ .

$$f(n) = (n-1)f(n-1) = (n-1)(n-2)f(n-2) = \dots = (n-1)!f(1)$$

$$U = \frac{f(11)}{(11-1)f(11-3)} = \frac{10!f(1)}{10 \times 7!f(1)} = 8 \times 9 = 72$$

I2.1 Let  $[x]$  be the largest integer not greater than  $x$ . If  $a = \left[ (\sqrt{3} - \sqrt{2})^{2009} \right] + 16$ , find the value of  $a$ .

$$0 < \sqrt{3} - \sqrt{2} = \frac{1}{\sqrt{3} + \sqrt{2}} < 1$$

$$0 < (\sqrt{3} - \sqrt{2})^{2009} < 1 \Rightarrow a = \left[ (\sqrt{3} - \sqrt{2})^{2009} \right] + 16 = 0 + 16 = 16$$

I2.2 In the coordinate plane, if the area of the triangle formed by the  $x$ -axis,  $y$ -axis and the line  $3x + ay = 12$  is  $b$  square units, find the value of  $b$ .

$$3x + 16y = 12; \text{ x-intercept} = 4, \text{ y-intercept} = \frac{3}{4}$$

$$\text{area} = b = \frac{1}{2} \cdot 4 \cdot \frac{3}{4} = \frac{3}{2}$$

I2.3 Given that  $x - \frac{1}{x} = 2b$  and  $x^3 - \frac{1}{x^3} = c$ , find the value of  $c$ .

$$x - \frac{1}{x} = 3 \Rightarrow x^2 - 2 + \frac{1}{x^2} = 9 \Rightarrow x^2 + \frac{1}{x^2} = 11$$

$$c = x^3 - \frac{1}{x^3} = \left( x - \frac{1}{x} \right) \left( x^2 + 1 + \frac{1}{x^2} \right) = 3 \times (11 + 1) = 36$$

I2.4 In Figure 1,  $\alpha = c$ ,  $\beta = 43$ ,  $\gamma = 59$  and  $\omega = d$ , find the value of  $d$ .


$\angle BAC = \omega^\circ$  ( $\angle$ s in the same seg.)

$\angle ACD = \beta^\circ$  ( $\angle$ s in the same seg.)

$\angle BAD + \angle BCD = 180^\circ$  (opp.  $\angle$ s cyclic quad.)

$$c + d + 43 + 59 = 180$$

$$d = 180 - 43 - 59 - 36 = 42 \quad (\because c = 36)$$



13.1 Given that  $\frac{4}{\sqrt{6} + \sqrt{2}} - \frac{1}{\sqrt{3} + \sqrt{2}} = \sqrt{a} - \sqrt{b}$ . If  $m = a - b$ , find the value of  $m$ .

$$\begin{aligned} \frac{4}{\sqrt{6} + \sqrt{2}} - \frac{1}{\sqrt{3} + \sqrt{2}} &= \frac{4}{\sqrt{6} + \sqrt{2}} \cdot \frac{\sqrt{6} - \sqrt{2}}{\sqrt{6} - \sqrt{2}} - \frac{1}{\sqrt{3} + \sqrt{2}} \cdot \frac{\sqrt{3} - \sqrt{2}}{\sqrt{3} - \sqrt{2}} \\ &= \frac{4(\sqrt{6} - \sqrt{2})}{6 - 2} - \frac{\sqrt{3} - \sqrt{2}}{3 - 2} \\ &= \sqrt{6} - \sqrt{2} - (\sqrt{3} - \sqrt{2}) \\ &= \sqrt{6} - \sqrt{3} \end{aligned}$$

$$a = 6, b = 3; m = 6 - 3 = 3$$

13.2 In figure 1,  $PQR$  is a right-angled triangle and  $RSTU$  is a rectangle. Let  $A$ ,  $B$  and  $C$  be the areas of the corresponding regions. If  $A : B = m : 2$  and  $A : C = n : 1$ , find the value of  $n$ .

$$A : B = 3 : 2, A : C = n : 1 \Rightarrow A : B : C = 3n : 2n : 3$$

$$\text{Let } SR = TU = x, QU = y$$

$\Delta PTS \sim \Delta TQU \sim \Delta PQR$  (equiangular)

$$S_{\Delta PTS} : S_{\Delta TQU} : S_{\Delta PQR} = A : C : (A + B + C) = 3n : 3 : (5n + 3)$$

$$x^2 : y^2 : (x + y)^2 = 3n : 3 : (5n + 3)$$

$$\frac{y}{x} = \frac{1}{\sqrt{n}} \dots (1), \quad \frac{x + y}{y} = \frac{\sqrt{5n + 3}}{\sqrt{3}} \dots (2)$$

$$\text{From (2): } \frac{x}{y} + 1 = \frac{\sqrt{5n + 3}}{\sqrt{3}} \dots (3)$$


$$\text{Sub. (1) into (3): } \sqrt{n} + 1 = \frac{\sqrt{5n + 3}}{\sqrt{3}}$$

$$\sqrt{3n} + \sqrt{3} = \sqrt{5n + 3}$$

$$(\sqrt{3n} + \sqrt{3})^2 = 5n + 3$$

$$3n + 6\sqrt{n} + 3 = 5n + 3$$

$$6\sqrt{n} = 2n \Rightarrow n = 9$$



Method 2

$$\text{Let } SR = x, PS = z$$

Join  $TR$  which bisects the area of the rectangle.

$$\frac{A}{B} = \frac{3}{2} \Rightarrow \frac{A}{\frac{B}{2}} = \frac{3}{1}$$

$$\frac{S_{\Delta TPS}}{S_{\Delta TSR}} = \frac{3}{1} \Rightarrow \frac{z}{x} = \frac{3}{1} \dots (4)$$

$\therefore \Delta PTS \sim \Delta TQU$  (equiangular)

$$\therefore \frac{A}{C} = \frac{n}{1} \Rightarrow \left(\frac{z}{x}\right)^2 = \frac{n}{1}$$

$$n = 3^2 = 9 \text{ (by (4))}$$

13.3 Let  $x_1, x_2, x_3, x_4$  be real numbers and  $x_1 \neq x_2$ . If  $(x_1 + x_3)(x_1 + x_4) = (x_2 + x_3)(x_2 + x_4) = n - 10$  and  $p = (x_1 + x_3)(x_2 + x_3) + (x_1 + x_4)(x_2 + x_4)$ , find the value of  $p$ .

$$\text{Reference: 2001-02 Heat Individual Q7: If } \frac{(a-b)(c-d)}{(b-c)(d-a)} = 3, \text{ find } \dots \frac{(a-c)(b-d)}{(a-b)(c-d)}$$

$$x_1^2 + x_1x_3 + x_1x_4 + x_3x_4 = x_2^2 + x_2x_3 + x_2x_4 + x_3x_4 = -1$$

$$x_1^2 - x_2^2 + x_1x_3 - x_2x_3 + x_1x_4 - x_2x_4 = 0$$

$$(x_1 - x_2)(x_1 + x_2 + x_3 + x_4) = 0 \Rightarrow x_1 + x_2 + x_3 + x_4 = 0 \dots (1)$$

$$p = (x_1 + x_3)(x_2 + x_3) + (x_1 + x_4)(x_2 + x_4)$$

$$= -(x_2 + x_3)(x_2 + x_4) - (x_1 + x_3)(x_1 + x_4) \text{ (by (1), } x_1 + x_3 = -(x_2 + x_4), x_2 + x_4 = -(x_1 + x_3))$$

$$= 1 + 1 = 2 \text{ (given } (x_1 + x_3)(x_1 + x_4) = (x_2 + x_3)(x_2 + x_4) = -1)$$

13.4 The total number of students in a school is a multiple of 7 and not less than 1000. Given that the same remainder 1 will be obtained when the number of students is divided by  $p + 1$ ,  $p + 2$  and  $p + 3$ . Let  $q$  be the least of the possible numbers of students in the school, find  $q$ .

$$p + 1 = 3, p + 2 = 4, p + 3 = 5; \text{ HCF} = 1, \text{ LCM} = 60$$

$$q = 60m + 1, \text{ where } m \text{ is an integer.}$$

$$\therefore q \geq 1000 \text{ and } q = 7k, k \text{ is an integer.}$$

$$60m + 1 = 7k$$

$$7k - 60m = 1$$


$$k = 43, m = 5 \text{ satisfies the equation}$$

$$k = 43 + 60t, 7k \geq 1000 \Rightarrow 7(43 + 60t) \geq 1000 \Rightarrow t \geq 2 \Rightarrow \text{Least } q = 7 \times (43 + 60 \times 2) = 1141$$

I4.1 Given that  $x_0^2 + x_0 - 1 = 0$ . If  $m = x_0^3 + 2x_0^2 + 2$ , find the value of  $m$ .

$$m = x_0^3 + 2x_0^2 + 2 = x_0^3 + x_0^2 - x_0 + x_0^2 + x_0 - 1 + 3 = 3$$

I4.2 In Figure 1,  $\triangle BAC$  is a right-angled triangle,  $AB = AC = m$  cm. Suppose that the circle with diameter  $AB$  intersects the line  $BC$  at  $D$ , and the total area of the shaded region is  $n$  cm<sup>2</sup>. Find the value of  $n$ .



$$AB = AC = 3 \text{ cm}; \angle ADB = 90^\circ (\angle \text{ in semi-circle})$$

$$\text{Shaded area} = \text{area of } \triangle ACD = \frac{1}{2} \text{ area of } \triangle ABC = \frac{1}{2} \cdot \frac{1}{2} \cdot 3 \cdot 3 \text{ cm}^2$$

$$n = \frac{9}{4}$$

I4.3 Given that  $p = 4n \left( \frac{1}{2^{2009}} \right)^{\log(1)}$ , find the value of  $p$ .

$$p = 4n \left( \frac{1}{2^{2009}} \right)^0 = 4 \cdot \frac{9}{4} = 9$$

I4.4 Let  $x$  and  $y$  be real numbers satisfying the equation  $(x - \sqrt{p})^2 + (y - \sqrt{p})^2 = 1$ . If  $k = \frac{y}{x - 3}$

and  $q$  is the least possible values of  $k^2$ , find the value of  $q$ .

$$(x - 3)^2 + (y - 3)^2 = 1 \dots (1)$$

$$\text{Sub. } y = k(x - 3) \text{ into (1): } (x - 3)^2 + [k(x - 3) - 3]^2 = 1$$

$$(x - 3)^2 + k^2(x - 3)^2 - 6k(x - 3) + 9 = 1$$

$$(1 + k^2)(x - 3)^2 - 6k(x - 3) + 8 = 0 \Rightarrow (1 + k^2)t^2 - 6kt + 8 = 0; \text{ where } t = x - 3$$

$$\text{For real values of } t: \Delta = 4[3^2k^2 - 8(1 + k^2)] \geq 0$$

$$k^2 \geq 8$$

The least possible value of  $k^2 = q = 8$ .

Method 2

The circle  $(x - 3)^2 + (y - 3)^2 = 1$  intersects with the variable line  $y = k(x - 3)$  which passes through a fixed point  $(3, 0)$ , where  $k$  is the slope of the line.

From the graph, the extreme points when the variable line touches the circle at  $B$  and  $C$ .

$H(3, 3)$  is the centre of the circle.

In  $\triangle ABH$ ,  $\angle ABH = 90^\circ$  ( $\angle$  in semi-circle)


Let  $\angle HAB = \theta$ ,  $AH = 3$ ,  $HB = 1$ ,

$$AB = \sqrt{8} = 2\sqrt{2} \text{ (Pyth. thm)}$$

$$\tan \theta = \frac{1}{\sqrt{8}}$$

$$\text{Slope of } AB = \tan(90^\circ - \theta) = \sqrt{8}$$

$$\text{Least possible } k^2 = q = 8$$



GS.1 In Figure 1,  $BD$ ,  $FC$ ,  $GC$  and  $FE$  are straight lines.

If  $z = a + b + c + d + e + f + g$ , find the value of  $z$ .

$$a^\circ + b^\circ + g^\circ + \angle BHG = 360^\circ \quad (\angle\text{s sum of polygon } ABHG)$$


$$c^\circ + f^\circ = \angle CJE \quad (\text{ext. } \angle \text{ of } \triangle CFJ)$$

$$c^\circ + f^\circ + e^\circ + d^\circ + \angle JHD = 360^\circ \quad (\angle\text{s sum of polygon } JHDE)$$

$$a^\circ + b^\circ + g^\circ + \angle BHG + c^\circ + f^\circ + e^\circ + d^\circ + \angle JHD = 720^\circ$$

$$a^\circ + b^\circ + c^\circ + d^\circ + e^\circ + f^\circ + g^\circ + 180^\circ = 720^\circ$$

$$z = 540$$



GS.2 If  $R$  is the remainder of  $1^6 + 2^6 + 3^6 + 4^6 + 5^6 + 6^6$  divided by 7, find the value of  $R$ .

$$x^6 + y^6 = (x + y)(x^5 - x^4y + x^3y^2 - x^2y^3 + xy^4 - y^5) + 2y^6$$

$$6^6 + 1^6 = 7Q_1 + 2; \quad 5^6 + 2^6 = 7Q_2 + 2 \times 2^6; \quad 4^6 + 3^6 = 7Q_3 + 2 \times 3^6$$

$$2 + 2 \times 2^6 + 2 \times 3^6 = 2(1 + 64 + 729) = 1588 = 7 \times 226 + 6; \quad R = 6$$

$$\begin{aligned} \text{Method 2} \quad 1^6 + 2^6 + 3^6 + 4^6 + 5^6 + 6^6 &\equiv 1^6 + 2^6 + 3^6 + (-3)^6 + (-2)^6 + (-1)^6 \pmod{7} \\ &\equiv 2(1^6 + 2^6 + 3^6) \pmod{7} \\ &\equiv 2(1 + 64 + 729) \pmod{7} \\ &\equiv 2(1 + 1 + 1) \pmod{7} \\ &\equiv 6 \pmod{7} \end{aligned}$$

GS.3 If  $14!$  is divisible by  $6^k$ , where  $k$  is an integer, find the largest possible value of  $k$ .

We count the number of factors of 3 in  $14!$ . They are 3, 6, 9, 12. So there are 5 factors of 3.

$$k = 5$$

GS.4 Let  $x$ ,  $y$  and  $z$  be real numbers that satisfy  $x + \frac{1}{y} = 4$ ,  $y + \frac{1}{z} = 1$  and  $z + \frac{1}{x} = \frac{7}{3}$ . Find the value of

$xyz$ .

Method 1

$$\text{From (1), } x = 4 - \frac{1}{y} = \frac{4y-1}{y}$$

$$\Rightarrow \frac{1}{x} = \frac{y}{4y-1} \quad \dots\dots (4)$$

$$\text{Sub. (4) into (3): } z + \frac{y}{4y-1} = \frac{7}{3}$$

$$z = \frac{7}{3} - \frac{y}{4y-1} \quad \dots\dots (5)$$

$$\text{From (2): } \frac{1}{z} = 1 - y$$

$$z = \frac{1}{1-y} \quad \dots\dots (6)$$

$$(5) = (6): \frac{1}{1-y} = \frac{7}{3} - \frac{y}{4y-1}$$

$$\frac{1}{1-y} = \frac{28y-7-3y}{3(4y-1)}$$

$$3(4y-1) = (1-y)(25y-7)$$

$$12y-3 = -25y^2-7+32y$$

$$25y^2-20y+4=0$$

$$(5y-2)^2=0$$

$$y = \frac{2}{5}$$

$$\text{Sub. } y = \frac{2}{5} \text{ into (6): } z = \frac{1}{1-\frac{2}{5}} = \frac{5}{3}$$

$$\text{Sub. } y = \frac{2}{5} \text{ into (1): } x + \frac{5}{2} = 4 \Rightarrow x = \frac{3}{2}$$

$$xyz = \frac{2}{5} \times \frac{5}{3} \times \frac{3}{2} = 1$$

Method 2

$$\begin{cases} x + \frac{1}{y} = 4 \dots\dots (1) \\ y + \frac{1}{z} = 1 \dots\dots (2) \\ z + \frac{1}{x} = \frac{7}{3} \dots\dots (3) \end{cases}$$

$$\left\{ \begin{aligned} (1) \times (2): \quad xy + 1 + \frac{x}{z} + \frac{1}{yz} &= 4 \\ x \left( y + \frac{1}{z} \right) + \frac{1}{yz} &= 3 \end{aligned} \right.$$

$$\text{Sub. (2) into the eqt.: } x + \frac{x}{xyz} = 3$$

$$\text{Let } a = xyz, \text{ then } x + \frac{x}{a} = 3 \dots\dots (4)$$

$$(2) \times (3): y \left( \frac{7}{3} \right) + \frac{y}{a} = \frac{4}{3} \Rightarrow y \left( \frac{7}{3} + \frac{1}{a} \right) = \frac{4}{3} \dots\dots (5)$$

$$(1) \times (3): z(4) + \frac{z}{a} = \frac{25}{3} \Rightarrow z \left( 4 + \frac{1}{a} \right) = \frac{25}{3} \dots\dots (6)$$

$$(4) \times (5) \times (6): a \left( 1 + \frac{1}{a} \right) \left( \frac{7}{3} + \frac{1}{a} \right) \left( 4 + \frac{1}{a} \right) = \frac{100}{3}$$

$$\frac{(a+1)(7a+3)(4a+1)}{3a^2} = \frac{100}{3}$$

$$\text{which reduces to } 28a^3 - 53a^2 + 22a + 3 = 0$$

$$\Rightarrow (a-1)^2(28a+3) = 0$$

$$\therefore a = 1$$

G1.1 Given some triangles with side lengths  $a$  cm, 2 cm and  $b$  cm, where  $a$  and  $b$  are integers and  $a \leq 2 \leq b$ . If there are  $q$  non-congruent classes of triangles satisfying the above conditions, find the value of  $q$ .

When  $a = 1$ , possible  $b = 2$

When  $a = 2$ , possible  $b = 2$  or 3

$\therefore q = 3$

G1.2 Given that the equation  $|x| - \frac{4}{x} = \frac{3|x|}{x}$  has  $k$  distinct real root(s), find the value of  $k$ .

When  $x > 0$ :  $x^2 - 4 = 3x \Rightarrow x^2 - 3x - 4 = 0 \Rightarrow (x + 1)(x - 4) = 0 \Rightarrow x = 4$

When  $x < 0$ :  $-x^2 - 4 = -3x \Rightarrow x^2 - 3x + 4 = 0$ ;  $D = 9 - 16 < 0 \Rightarrow$  no real roots.

$k = 1$  (There is only one real root.)

G1.3 Given that  $x$  and  $y$  are non-zero real numbers satisfying the equations  $\frac{\sqrt{x}}{\sqrt{y}} - \frac{\sqrt{y}}{\sqrt{x}} = \frac{7}{12}$  and

$x - y = 7$ . If  $w = x + y$ , find the value of  $w$ .

The first equation is equivalent to  $\frac{x - y}{\sqrt{xy}} = \frac{7}{12} \Rightarrow \sqrt{xy} = 12 \Rightarrow xy = 144$

Sub.  $y = \frac{144}{x}$  into  $x - y = 7$ :  $x - \frac{144}{x} = 7 \Rightarrow x^2 - 7x - 144 = 0 \Rightarrow (x + 9)(x - 16) = 0$

$x = -9$  or 16; when  $x = -9$ ,  $y = -16$  (rejected  $\because \sqrt{x}$  is undefined); when  $x = 16$ ;  $y = 9$

$w = 16 + 9 = 25$

G1.4 Given that  $x$  and  $y$  are real numbers and  $\left|x - \frac{1}{2}\right| + \sqrt{y^2 - 1} = 0$ . Let  $p = |x| + |y|$ , find the value of  $p$ .

Reference 2006 Final Individual 4.2 ...  $y^2 + 4y + 4 + \sqrt{x + y + k} = 0$ . If  $r = |xy|$ , ...

Both  $\left|x - \frac{1}{2}\right|$  and  $\sqrt{y^2 - 1}$  are non-negative numbers.

The sum of two non-negative numbers = 0 means each of them is zero

$x = \frac{1}{2}$ ,  $y = \pm 1$ ;  $p = \frac{1}{2} + 1 = \frac{3}{2}$

G2.1 Given  $\tan \theta = \frac{5}{12}$ , where  $180^\circ \leq \theta \leq 270^\circ$ . If  $A = \cos \theta + \sin \theta$ , find the value of  $A$ .

$$\cos \theta = -\frac{12}{13}, \sin \theta = -\frac{5}{13}$$

$$A = -\frac{12}{13} - \frac{5}{13} = -\frac{17}{13}$$

G2.2 Let  $[x]$  be the largest integer not greater than  $x$ . If  $B = \left[ 10 + \sqrt{10 + \sqrt{10 + \sqrt{10 + \dots}}} \right]$ , find the value of  $B$ .

Reference 2007 Final Group 2.2 ...  $x \geq 3 + \sqrt{3 + \sqrt{3 + \sqrt{3 + \sqrt{3 + \sqrt{3}}}}}$  ...

$$\text{Let } y = \sqrt{10 + \sqrt{10 + \sqrt{10 + \dots}}}$$

$$y^2 = 10 + \sqrt{10 + \sqrt{10 + \sqrt{10 + \dots}}} = 10 + y$$

$$y^2 - y - 10 = 0$$

$$y = \frac{1 + \sqrt{41}}{2} \text{ or } \frac{1 - \sqrt{41}}{2} \text{ (rejected)}$$

$$6 < \sqrt{41} < 7 \Rightarrow \frac{7}{2} < \frac{1 + \sqrt{41}}{2} < 4$$

$$13.5 < 10 + \sqrt{10 + \sqrt{10 + \sqrt{10 + \dots}}} < 14; B = 13$$

G2.3 Let  $a \oplus b = ab + 10$ . If  $C = (1 \oplus 2) \oplus 3$ , find the value of  $C$ .

$$1 \oplus 2 = 2 + 10 = 12; C = 12 \oplus 3 = 36 + 10 = 46$$


G2.4 In the coordinate plane, the area of the region bounded by the following lines is  $D$  square units, find the value of  $D$ .

$$L_1: y - 2 = 0$$

$$L_2: y + 2 = 0$$

$$L_3: 4x + 7y - 10 = 0$$

$$L_4: 4x + 7y + 20 = 0$$



It is easy to show that the bounded region is a parallelogram  $ABCD$  with vertices  $A(-1.5, 2)$ ,  $B(6, 2)$ ,  $C(-1, 2)$ ,  $D(-8.5, 2)$ .

$$\text{The area } D = |6 - (-1.5)| \times |2 - (-2)| = 7.5 \times 4 = 30$$

Group Event 3

G3.1 Let  $[x]$  be the largest integer not greater than  $x$ .

If  $A = \left\lceil \frac{2008 \times 80 + 2009 \times 130 + 2010 \times 180}{2008 \times 15 + 2009 \times 25 + 2010 \times 35} \right\rceil$ , find the value of  $A$ .

Reference 2008 Final Group Spare 4 Calculate the value of  $\frac{2008^3 + 4015^3}{2007^3 + 4015^3}$ .

Let  $a = 2009$ ,  $b = 130$ ,  $c = 25$

$$\begin{aligned} \frac{2008 \times 80 + 2009 \times 130 + 2010 \times 180}{2008 \times 15 + 2009 \times 25 + 2010 \times 35} &= \frac{(a-1)(b-50) + ab + (a+1)(b+50)}{(a-1)(c-10) + ac + (a+1)(c+10)} \\ &= \frac{ab - b - 50a + 50 + ab + ab + b + 50a + 50}{ac - c - 10a + 10 + ac + ac + c + 10a + 10} \\ &= \frac{3ab + 100}{3ac + 20} = \frac{3 \cdot 2009 \cdot 130 + 100}{3 \cdot 2009 \cdot 25 + 20} = \frac{783610}{150695} = 5 + d \end{aligned}$$

where  $0 < d < 1$ ;  $A = 5$

G3.2 There are  $R$  zeros at the end of  $\underbrace{99\dots9}_{2009 \text{ of } 9\text{'s}} \times \underbrace{99\dots9}_{2009 \text{ of } 9\text{'s}} + 1 \underbrace{99\dots9}_{2009 \text{ of } 9\text{'s}}$ , find the value of  $R$ .

$$\begin{aligned} \underbrace{99\dots9}_{2009 \text{ of } 9\text{'s}} \times \underbrace{99\dots9}_{2009 \text{ of } 9\text{'s}} + 1 \underbrace{99\dots9}_{2009 \text{ of } 9\text{'s}} &= \left( \begin{matrix} 1 & \underbrace{0\dots0}_{2009 \text{ of } 0\text{'s}} & -1 \end{matrix} \right) \times \left( \begin{matrix} 1 & \underbrace{0\dots0}_{2009 \text{ of } 0\text{'s}} & -1 \end{matrix} \right) + \left( \begin{matrix} 2 & \underbrace{0\dots0}_{2009 \text{ of } 0\text{'s}} & -1 \end{matrix} \right) \\ &= (10^{2009} - 1)(10^{2009} - 1) + 2 \times 10^{2009} - 1 \\ &= 10^{4018} - 2 \times 10^{2009} + 1 + 2 \times 10^{2009} - 1 = 10^{4018} \end{aligned}$$


$R = 4018$

G3.3 In Figure 1, a square of side length  $Q$  cm is inscribed in a semi-circle of radius 2 cm. Find the value of  $Q$ .

$$Q^2 + \left(\frac{Q}{2}\right)^2 = 4 \quad (\text{Pyth. Thm})$$

$$5Q^2 = 16$$

$$Q = \frac{4}{\sqrt{5}} = \frac{4\sqrt{5}}{5}$$



G3.4 In Figure 2, the sector  $OAB$  has radius 4 cm and  $\angle AOB$  is a right angle. Let the semi-circle with diameter  $OB$  be centred at  $I$  with  $IJ \parallel OA$ , and  $IJ$  intersects the semi-circle at  $K$ . If the area of the shaded region is  $T$  cm<sup>2</sup>, find the value of  $T$ . (Take  $\pi = 3$ )


$$OI = 2 \text{ cm}, OJ = 4 \text{ cm}, \cos \angle IOJ = \frac{OI}{OJ} = \frac{1}{2}; \angle IOJ = \frac{\pi}{3}$$

$$S_{BLJ} = S_{\text{sector } OBJ} - S_{\Delta OIJ} = \left( \frac{1}{2} \cdot 4^2 \cdot \frac{\pi}{3} - \frac{1}{2} \cdot 2 \cdot 4 \sin 60^\circ \right) \text{cm}^2$$

$$= \left( \frac{8\pi}{3} - 2\sqrt{3} \right) \text{cm}^2$$

$$\text{Shaded area} = S_{BLJ} - S_{BIK} = \frac{8\pi}{3} - 2\sqrt{3} - \frac{1}{4} \pi \cdot 2^2 = \frac{5\pi}{3} - 2\sqrt{3} \text{ cm}^2$$


$$T = 5 - 2\sqrt{3}$$



G4.1 Let  $P$  be a real number. If  $\sqrt{3-2P} + \sqrt{1-2P} = 2$ , find the value of  $P$ .

$$\begin{aligned} (\sqrt{3-2P})^2 &= (2 - \sqrt{1-2P})^2 \\ 3 - 2P &= 4 - 4\sqrt{1-2P} + 1 - 2P \\ 4\sqrt{1-2P} &= 2 \\ 4(1-2P) &= 1 \\ P &= \frac{3}{8} \end{aligned}$$

G4.2 In Figure 1, let  $AB$ ,  $AC$  and  $BC$  be the diameters of the corresponding three semi-circles. If  $AC = BC = 1$  cm and the area of the shaded region is  $R$  cm<sup>2</sup>. Find the value of  $R$ .




Reference 1994 Heat Individual Q9

$$AB = \sqrt{2}$$

$$\text{Shaded area} = R \text{ cm}^2 = S_{\text{circle with diameter } AC} - 2 S_{\text{segment } AC}$$

$$R = \pi \left( \frac{1}{2} \right)^2 - \left[ \frac{1}{2} \pi \cdot \left( \frac{\sqrt{2}}{2} \right)^2 - \frac{1}{2} \cdot 1^2 \right] = \frac{1}{2}$$

G4.3 In Figure 2,  $AC$ ,  $AD$ ,  $BD$ ,  $BE$  and  $CF$  are straight lines. If  $\angle A + \angle B + \angle C + \angle D = 140^\circ$  and  $a + b + c = S$ , find the value of  $S$ .



$$\angle CFD = \angle A + \angle C \text{ (ext. } \angle \text{ of } \Delta)$$

$$\angle AEB = \angle B + \angle D \text{ (ext. } \angle \text{ of } \Delta)$$

$$a^\circ = \angle A + \angle AEB = \angle A + \angle B + \angle D \text{ (ext. } \angle \text{ of } \Delta)$$

$$c^\circ = \angle D + \angle CFD = \angle D + \angle A + \angle C \text{ (ext. } \angle \text{ of } \Delta)$$

$$b^\circ + \angle A + \angle D = 180^\circ \text{ (} \angle \text{s sum of } \Delta)$$

$$\begin{aligned} a^\circ + b^\circ + c^\circ &= \angle A + \angle B + \angle D + 180^\circ - (\angle A + \angle D) \\ &\quad + \angle D + \angle A + \angle C \\ &= \angle A + \angle B + \angle C + \angle D + 180^\circ \end{aligned}$$

$$S = a + b + c = 140 + 180 = 320$$

G4.4 Let  $Q = \log_{2+\sqrt{2^2-1}}(2 - \sqrt{2^2-1})$ , find the value of  $Q$ .

$$Q = \frac{\log(2 - \sqrt{3})}{\log(2 + \sqrt{3})} = \frac{\log(2 - \sqrt{3})}{\log(2 + \sqrt{3}) \cdot \frac{(2 - \sqrt{3})}{(2 - \sqrt{3})}} = \frac{\log(2 - \sqrt{3})}{\log \frac{1}{2 - \sqrt{3}}} = \frac{\log(2 - \sqrt{3})}{-\log(2 - \sqrt{3})} = -1$$

$$\text{method 2 } Q = \log_{2+\sqrt{3}}(2 - \sqrt{3}) \cdot \frac{(2 + \sqrt{3})}{(2 + \sqrt{3})} = \log_{2+\sqrt{3}} \frac{1}{(2 + \sqrt{3})} = \log_{2+\sqrt{3}}(2 + \sqrt{3})^{-1} = -1$$