

Laparoscopic Cystogastrostomy for Management of Pancreatic Congenital Cyst

Poschong Suesat M.D.

Pancreatic cysts

Pseudocysts

Congenital cysts

Cystic neoplasms

Pseudocysts

Causes - pancreatitis, blunt trauma **Symptoms** - persistent abdominal pain, abdominal mass, norsia and vomitting Complications - infected cyst, ruptured

Congenital cysts

- True cyst
- Epithelial lining (cuboidal)
- Von Hippel-Lindau disease (cystic lesions :- brain, retina, adrenal glands, kidneys, pancreas)

Cystic neoplasms

1. Mucinous cystadenoma

(columnar, papillary mucin-producing epithelium)

2. Mucinous duct ectasia, IPMN

(pancreatic ducts, mucin-producing columnar epithelium papillary projection)

3. Serous cystadenoma

(glycogen-rich cytoplasm cuboidal cells)

4. Papillary cystic tumor

(solid and pseudopapillary neoplasm)

Treatments of cystic neoplasms

Observation

(asymptomatic precancerous cysts, Serous cystadenoma)

Distal pancreatectomy

(pancreatic tail)

Pancreaticoduodenectomy

(pancreatic head)

Treatments of pseudocysts

Observation

(asymptomatic)

Resection

(pancreatic tail)


External drainage

(thin wall, rupture, fistula, recurrent cysts)


Internal drainage


(cystojejunostomy, cystogastrostomy, cystoduodenostomy)

Open conventional cystogastrostomy












Laparoscopic surgery

cholecystectomy fundoplication splenectomy adrenalectomy

Congenital cyst, a case report



Laparoscopic cystogastrostomy





Laparoscopic cystogastrostomy





#